



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 11, 2023 – 02:55 pm BST

PDB ID : 7ZD1  
Title : Crystal structure of Pseudomonas aeruginosa S-adenosyl-L-homocysteine hydrolase inhibited by Hg<sup>2+</sup> ions  
Authors : Malecki, P.H.; Gawel, M.; Brzezinski, K.  
Deposited on : 2022-03-29  
Resolution : 1.56 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.32.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	472	91% 6% ..
1	B	472	90% 7% .
1	C	472	91% 6% .
1	D	472	% 91% 6% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
7	HG	A	509	-	-	X	-
7	HG	C	508	-	-	X	-

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 16750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosylhomocysteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3592	2269	619	681	23	0	7	0
1	B	460	3573	2253	618	680	22	0	4	0
1	C	461	3603	2275	620	685	23	0	9	0
1	D	461	3624	2287	627	687	23	0	10	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9I685
A	-1	ASN	-	expression tag	UNP Q9I685
A	0	ALA	-	expression tag	UNP Q9I685
B	-2	SER	-	expression tag	UNP Q9I685
B	-1	ASN	-	expression tag	UNP Q9I685
B	0	ALA	-	expression tag	UNP Q9I685
C	-2	SER	-	expression tag	UNP Q9I685
C	-1	ASN	-	expression tag	UNP Q9I685
C	0	ALA	-	expression tag	UNP Q9I685
D	-2	SER	-	expression tag	UNP Q9I685
D	-1	ASN	-	expression tag	UNP Q9I685
D	0	ALA	-	expression tag	UNP Q9I685

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
			Total	C	N	O			P	
2	A	1	Total	44	21	7	14	2	0	0
2	B	1	Total	44	21	7	14	2	0	0
2	C	1	Total	44	21	7	14	2	0	0
2	D	1	Total	44	21	7	14	2	0	0

- Molecule 3 is ADENOSINE (three-letter code: ADN) (formula:  $C_{10}H_{13}N_5O_4$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			19	10	5	4		
3	B	1	Total	C	N	O	0	0
			19	10	5	4		
3	C	1	Total	C	N	O	0	0
			19	10	5	4		
3	D	1	Total	C	N	O	0	0
			19	10	5	4		

- Molecule 4 is 1,4-BUTANEDIOL (three-letter code: BU1) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C O	0	0
			6	4 2		
4	D	1	Total	C O	0	0
			6	4 2		

- Molecule 5 is 1,3-PROPANDIOL (three-letter code: PDO) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 5 3 2	0	0
5	A	1	Total C O 5 3 2	0	0
5	B	1	Total C O 5 3 2	0	0
5	B	1	Total C O 5 3 2	0	0
5	C	1	Total C O 5 3 2	0	0
5	C	1	Total C O 5 3 2	0	0
5	C	1	Total C O 5 3 2	0	0

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total K 1 1	0	0
6	B	1	Total K 1 1	0	0
6	C	1	Total K 1 1	0	0
6	D	1	Total K 1 1	0	0

- Molecule 7 is MERCURY (II) ION (three-letter code: HG) (formula: Hg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	5	Total Hg 5 5	0	1
7	B	5	Total Hg 5 5	0	1
7	C	5	Total Hg 5 5	0	1
7	D	5	Total Hg 5 5	0	1

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	1	Total Cl 1 1	0	0

- Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	567	Total O 572 572	0	10
9	B	465	Total O 472 472	0	11
9	C	512	Total O 519 519	0	11
9	D	465	Total O 471 471	0	12




### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Adenosylhomocysteinase

Chain A:  91% 6% ..





- Molecule 1: Adenosylhomocysteinase

Chain B:  90% 7% .





- Molecule 1: Adenosylhomocysteinase

Chain C:  91% 6% .



- Molecule 1: Adenosylhomocysteinase

Chain D:  91% 6% .





## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	177.42Å 134.20Å 108.73Å 90.00° 105.95° 90.00°	Depositor
Resolution (Å)	105.47 – 1.56 105.47 – 1.56	Depositor EDS
% Data completeness (in resolution range)	99.5 (105.47-1.56) 99.5 (105.47-1.56)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.48 (at 1.56Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.123 , 0.179 0.134 , 0.196	Depositor DCC
$R_{free}$ test set	1033 reflections (0.30%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.6	Xtrriage
Anisotropy	0.196	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.000 for $-1/2^*h+1/2^*k+1, 1/2^*h-1/2^*k+1, 1/2^*h+1/2^*k$ 0.000 for $-1/2^*h-1/2^*k+1, -1/2^*h-1/2^*k-1, 1/2^*h-1/2^*k$	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	16750	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, BU1, CL, PDO, ADN, K, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.86	2/3670 (0.1%)	0.94	7/4962 (0.1%)
1	B	0.86	6/3645 (0.2%)	0.92	5/4928 (0.1%)
1	C	0.84	1/3687 (0.0%)	0.96	5/4985 (0.1%)
1	D	0.86	2/3701 (0.1%)	0.94	8/5000 (0.2%)
All	All	0.86	11/14703 (0.1%)	0.94	25/19875 (0.1%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	434	GLU	CD-OE2	8.81	1.35	1.25
1	D	434	GLU	CD-OE2	6.61	1.32	1.25
1	B	174	ASP	CG-OD1	6.00	1.39	1.25
1	A	79	GLU	CD-OE2	-5.86	1.19	1.25
1	D	355	ASP	C-O	5.48	1.33	1.23

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	241	ARG	NE-CZ-NH1	8.68	124.64	120.30
1	A	213	ARG	NE-CZ-NH2	-7.61	116.50	120.30
1	A	417	ARG	NE-CZ-NH1	7.25	123.93	120.30
1	C	277	ASP	CB-CG-OD2	-7.14	111.87	118.30
1	C	26	ARG	NE-CZ-NH2	-6.73	116.94	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3592	0	3613	20	0
1	B	3573	0	3582	18	0
1	C	3603	0	3623	21	0
1	D	3624	0	3632	31	0
2	A	44	0	26	1	0
2	B	44	0	26	1	0
2	C	44	0	26	1	0
2	D	44	0	26	1	0
3	A	19	0	13	2	0
3	B	19	0	13	2	0
3	C	19	0	13	2	0
3	D	19	0	13	3	0
4	A	6	0	10	1	0
4	D	6	0	10	2	0
5	A	10	0	16	0	0
5	B	10	0	16	0	0
5	C	15	0	24	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	5	0	0	2	0
7	B	5	0	0	2	0
7	C	5	0	0	3	0
7	D	5	0	0	2	0
8	D	1	0	0	0	0
9	A	572	0	0	6	0
9	B	472	0	0	4	0
9	C	519	0	0	1	0
9	D	471	0	0	9	0
All	All	16750	0	14682	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:85:CYS:HB3	7:C:508:HG:HG	0.94	1.11
1:A:119:CYS:SG	7:A:509:HG:HG	1.90	0.88
1:C:223[B]:GLN:HE22	1:C:290:LYS:HE3	1.38	0.85
1:C:119:CYS:SG	7:C:509:HG:HG	1.97	0.81
1:D:323[A]:HIS:O	1:D:373:LEU:HD11	1.81	0.79

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/472 (99%)	457 (98%)	9 (2%)	0	100	100
1	B	462/472 (98%)	451 (98%)	10 (2%)	1 (0%)	47	23
1	C	468/472 (99%)	456 (97%)	12 (3%)	0	100	100
1	D	469/472 (99%)	453 (97%)	15 (3%)	1 (0%)	47	23
All	All	1865/1888 (99%)	1817 (97%)	46 (2%)	2 (0%)	51	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	61	HIS
1	B	61	HIS

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/385 (100%)	376 (98%)	8 (2%)	53	24
1	B	381/385 (99%)	376 (99%)	5 (1%)	69	44
1	C	386/385 (100%)	383 (99%)	3 (1%)	81	66
1	D	386/385 (100%)	380 (98%)	6 (2%)	62	35
All	All	1537/1540 (100%)	1515 (99%)	22 (1%)	67	41

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	355	ASP
1	D	177	LYS
1	D	174	ASP
1	D	198	ASP
1	A	355	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	178	ASN
1	B	273	ASN
1	C	360	HIS
1	C	400	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 42 ligands modelled in this entry, 25 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	PDO	A	505	-	4,4,4	0.34	0	3,3,3	0.14	0
2	NAD	B	501	-	42,48,48	0.97	2 (4%)	50,73,73	0.99	2 (4%)
5	PDO	C	504	-	4,4,4	0.29	0	3,3,3	0.15	0
5	PDO	B	504	-	4,4,4	0.24	0	3,3,3	0.48	0
3	ADN	D	502	7	18,21,21	0.64	0	18,31,31	1.06	1 (5%)
2	NAD	C	501	-	42,48,48	1.04	4 (9%)	50,73,73	0.94	1 (2%)
5	PDO	A	504	-	4,4,4	0.53	0	3,3,3	0.45	0
4	BU1	A	503	-	5,5,5	0.50	0	4,4,4	0.54	0
4	BU1	D	503	-	5,5,5	0.32	0	4,4,4	0.20	0
2	NAD	D	501	-	42,48,48	1.15	3 (7%)	50,73,73	0.99	4 (8%)
3	ADN	B	502	-	18,21,21	1.00	1 (5%)	18,31,31	1.47	4 (22%)
5	PDO	C	505	-	4,4,4	0.32	0	3,3,3	0.31	0
2	NAD	A	501	-	42,48,48	1.04	3 (7%)	50,73,73	0.95	3 (6%)
3	ADN	C	502	7	18,21,21	0.96	0	18,31,31	1.29	1 (5%)
5	PDO	B	503	-	4,4,4	0.31	0	3,3,3	0.37	0
3	ADN	A	502	-	18,21,21	0.84	0	18,31,31	1.49	4 (22%)
5	PDO	C	503	-	4,4,4	0.27	0	3,3,3	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PDO	A	505	-	-	0/2/2/2	-
2	NAD	B	501	-	-	5/26/62/62	0/5/5/5
5	PDO	C	504	-	-	2/2/2/2	-
5	PDO	B	504	-	-	1/2/2/2	-
3	ADN	D	502	7	-	1/2/22/22	0/3/3/3
2	NAD	C	501	-	-	5/26/62/62	0/5/5/5
5	PDO	A	504	-	-	1/2/2/2	-
4	BU1	A	503	-	-	2/3/3/3	-

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BU1	D	503	-	-	2/3/3/3	-
2	NAD	D	501	-	-	5/26/62/62	0/5/5/5
3	ADN	B	502	-	-	0/2/22/22	0/3/3/3
5	PDO	C	505	-	-	1/2/2/2	-
2	NAD	A	501	-	-	5/26/62/62	0/5/5/5
3	ADN	C	502	7	-	0/2/22/22	0/3/3/3
5	PDO	B	503	-	-	0/2/2/2	-
3	ADN	A	502	-	-	0/2/22/22	0/3/3/3
5	PDO	C	503	-	-	1/2/2/2	-

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	NAD	O7N-C7N	4.40	1.32	1.24
2	C	501	NAD	C7N-N7N	3.48	1.39	1.33
2	D	501	NAD	C7N-N7N	3.44	1.39	1.33
2	A	501	NAD	C2B-C1B	-3.31	1.48	1.53
2	A	501	NAD	C3N-C7N	-3.19	1.45	1.50

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	ADN	C5-C6-N6	3.10	125.06	120.35
2	B	501	NAD	C2N-N1N-C1D	-2.96	112.55	119.14
3	B	502	ADN	O2'-C2'-C1'	2.95	121.75	110.85
3	B	502	ADN	C5-C6-N6	2.89	124.75	120.35
2	A	501	NAD	C5A-C6A-N6A	2.79	124.59	120.35

There are no chirality outliers.

5 of 31 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	NAD	O4D-C1D-N1N-C2N
2	A	501	NAD	O4D-C1D-N1N-C6N
2	A	501	NAD	C2D-C1D-N1N-C2N
2	A	501	NAD	C2D-C1D-N1N-C6N
2	B	501	NAD	O4D-C1D-N1N-C2N


There are no ring outliers.


11 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	NAD	1	0
3	D	502	ADN	3	0
2	C	501	NAD	1	0
4	A	503	BU1	1	0
4	D	503	BU1	2	0
2	D	501	NAD	1	0
3	B	502	ADN	2	0
2	A	501	NAD	1	0
3	C	502	ADN	2	0
3	A	502	ADN	2	0
5	C	503	PDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	461/472 (97%)	-0.55	1 (0%) 95 95	19, 26, 42, 96	0
1	B	460/472 (97%)	-0.44	2 (0%) 92 94	19, 31, 57, 83	0
1	C	461/472 (97%)	-0.56	1 (0%) 95 95	20, 29, 48, 90	0
1	D	461/472 (97%)	-0.42	4 (0%) 84 87	19, 30, 58, 87	0
All	All	1843/1888 (97%)	-0.49	8 (0%) 92 94	19, 28, 55, 96	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	10	PHE	4.6
1	D	10	PHE	4.6
1	B	409	LEU	4.2
1	A	355	ASP	4.0
1	D	324[A]	PHE	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	HG	D	509	1/1	0.68	0.10	72,72,72,72	1
4	BU1	D	503	6/6	0.72	0.10	62,70,80,83	0
5	PDO	C	504	5/5	0.84	0.12	62,64,77,78	0
5	PDO	A	504	5/5	0.84	0.10	47,59,68,69	0
7	HG	A	511[A]	1/1	0.86	0.41	69,69,69,69	1
7	HG	D	508[A]	1/1	0.87	0.21	67,67,67,67	1
5	PDO	B	503	5/5	0.89	0.10	52,56,66,67	0
5	PDO	A	505	5/5	0.89	0.14	55,57,66,81	0
5	PDO	C	505	5/5	0.89	0.11	55,68,72,79	0
5	PDO	C	503	5/5	0.91	0.10	49,53,70,77	0
5	PDO	B	504	5/5	0.91	0.08	52,55,70,79	0
7	HG	C	511	1/1	0.93	0.04	65,65,65,65	1
4	BU1	A	503	6/6	0.94	0.07	50,56,58,64	0
7	HG	B	510	1/1	0.95	0.10	69,69,69,69	1
3	ADN	B	502	19/19	0.95	0.12	22,24,52,52	0
3	ADN	C	502	19/19	0.95	0.12	20,22,52,53	0
3	ADN	A	502	19/19	0.95	0.12	19,21,48,49	0
7	HG	C	510[A]	1/1	0.97	0.17	65,65,65,65	1
3	ADN	D	502	19/19	0.97	0.08	21,25,35,35	0
7	HG	B	509[A]	1/1	0.97	0.17	39,39,39,39	1
2	NAD	B	501	44/44	0.97	0.06	19,23,26,28	0
2	NAD	D	501	44/44	0.98	0.06	20,22,26,26	0
2	NAD	A	501	44/44	0.98	0.05	18,21,23,26	0
2	NAD	C	501	44/44	0.98	0.06	22,24,26,28	0
7	HG	B	508	1/1	0.99	0.06	53,53,53,53	1
7	HG	A	508	1/1	0.99	0.05	36,36,36,36	1
7	HG	A	509	1/1	0.99	0.12	51,51,51,51	1
7	HG	C	507	1/1	0.99	0.05	35,35,35,35	1
7	HG	C	508	1/1	0.99	0.04	40,40,40,40	1
7	HG	C	509	1/1	0.99	0.11	55,55,55,55	1
7	HG	A	510	1/1	0.99	0.08	52,52,52,52	1
7	HG	A	507	1/1	0.99	0.03	27,27,27,27	1
7	HG	D	505	1/1	0.99	0.10	43,43,43,43	1
7	HG	D	506	1/1	0.99	0.03	38,38,38,38	1
7	HG	B	506	1/1	0.99	0.02	42,42,42,42	1
7	HG	B	507	1/1	0.99	0.07	42,42,42,42	1
6	K	D	504	1/1	1.00	0.07	25,25,25,25	0
6	K	A	506	1/1	1.00	0.09	22,22,22,22	0
7	HG	D	507	1/1	1.00	0.04	50,50,50,50	1
6	K	B	505	1/1	1.00	0.10	25,25,25,25	0
6	K	C	506	1/1	1.00	0.09	25,25,25,25	0
8	CL	D	510	1/1	1.00	0.10	28,28,28,28	1




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around HG A 511 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG D 508 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG C 511:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG B 510:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ADN B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ADN C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)













**Electron density around HG C 510 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ADN D 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG B 509 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAD B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NAD D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around NAD A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NAD C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG B 508:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HG A 508:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around HG A 509:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG C 507:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG C 508:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG C 509:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG A 510:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG A 507:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG D 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HG D 506:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around HG B 506:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HG B 507:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around K A 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HG D 507:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around K B 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.