



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2023 – 01:29 PM EDT

PDB ID : 1ZEA  
Title : Structure of the anti-cholera toxin antibody Fab fragment TE33 in complex with a D-peptide  
Authors : Scheerer, P.; Krauss, N.; Wessner, H.; Scholz, C.; Otte, L.; Seifert, M.; Kramer, A.; Schneider-Mergener, J.; Hoehne, W.  
Deposited on : 2005-04-18  
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	216	 6% 83% 16%
2	H	216	 2% 84% 11% ...
3	A	9	 22% 78%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DVA	A	1	-	-	X	-
3	DSN	A	9	-	-	-	X
4	CIT	L	1001	-	X	X	-
4	CIT	L	1002	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called monoclonal anti-cholera toxin IGG1 KAPPA antibody, L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	216	1678	1054	280	338	6	0	0	0


- Molecule 2 is a protein called monoclonal anti-cholera toxin IGG1 KAPPA antibody, H chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	209	1606	1030	260	309	7	0	0	0

- Molecule 3 is a protein called short synthetic D-amino acid peptide D2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	A	9	70	42	12	16	0	0	0

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			13	6	7		
4	L	1	Total	C	O	0	0
			13	6	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	136	Total	O	0	0
			136	136		
5	H	134	Total	O	0	0
			134	134		
5	A	6	Total	O	0	0
			6	6		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: monoclonal anti-cholera toxin IGG1 KAPPA antibody, L chain



- Molecule 2: monoclonal anti-cholera toxin IGG1 KAPPA antibody, H chain



- Molecule 3: short synthetic D-amino acid peptide D2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.64Å 108.86Å 40.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.00 – 1.78 22.84 – 1.78	Depositor EDS
% Data completeness (in resolution range)	93.4 (23.00-1.78) 93.4 (22.84-1.78)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.37 (at 1.78Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.198 , 0.248 0.190 , 0.196	Depositor DCC
$R_{free}$ test set	2041 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.2	Xtrriage
Anisotropy	0.166	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3656	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DVA, DTY, DGN, DHI, DAS, CIT, DSN, DPR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.57	0/1718	0.80	1/2328 (0.0%)
2	H	0.60	0/1654	0.86	2/2263 (0.1%)
3	A	0.59	0/3	1.64	0/2
All	All	0.59	0/3375	0.83	3/4593 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	177	LEU	CA-CB-CG	6.37	129.95	115.30
1	L	108	ARG	NE-CZ-NH2	-5.87	117.37	120.30
2	H	175	TYR	CB-CG-CD2	-5.03	117.98	121.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1678	0	1620	32	0
2	H	1606	0	1569	17	0
3	A	70	0	53	9	0
4	L	26	0	10	8	0
5	A	6	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	134	0	0	1	0
5	L	136	0	0	2	0
All	All	3656	0	3252	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:16:GLU:O	2:H:82(C):LEU:HD23	1.65	0.97
1:L:50:LYS:NZ	4:L:1001:CIT:H41	1.96	0.79
1:L:50:LYS:HD3	4:L:1001:CIT:O6	1.82	0.79
3:A:1:DVA:HB	3:A:2:DPR:HA	1.66	0.77
2:H:116:THR:HG21	5:H:287:HOH:O	1.85	0.76
2:H:136:VAL:HG23	2:H:185:SER:HA	1.77	0.67
1:L:108:ARG:HD3	1:L:109:ALA:O	1.96	0.66
1:L:50:LYS:HZ3	4:L:1001:CIT:H41	1.59	0.65
2:H:15:GLY:HA2	2:H:82(B):ASN:HD22	1.61	0.65
3:A:1:DVA:HA	3:A:3:GLY:H	1.67	0.60
1:L:50:LYS:HD3	4:L:1001:CIT:C6	2.32	0.59
1:L:80:ALA:HA	1:L:106:ILE:HD13	1.83	0.59
2:H:52(A):THR:H	3:A:5:DGN:HE22	1.50	0.57
1:L:49:TYR:HE2	4:L:1001:CIT:C6	2.18	0.57
1:L:144:ILE:HG22	1:L:163:TRP:CH2	2.40	0.56
3:A:1:DVA:HB	3:A:2:DPR:CA	2.36	0.56
1:L:49:TYR:HE2	4:L:1001:CIT:O6	1.87	0.55
1:L:80:ALA:HA	1:L:106:ILE:CD1	2.36	0.55
2:H:59:TYR:HB2	2:H:64:LYS:HG2	1.87	0.55
1:L:50:LYS:HZ2	4:L:1001:CIT:H41	1.69	0.55
1:L:189:HIS:O	1:L:211:ARG:HD3	2.08	0.53
2:H:143:LYS:HG2	2:H:176:THR:HG23	1.92	0.52
1:L:12:PRO:HA	1:L:105:GLU:O	2.11	0.51
1:L:107:LYS:HG2	1:L:140:TYR:OH	2.12	0.50
2:H:170:LEU:HB2	2:H:175:TYR:CE1	2.47	0.50
1:L:144:ILE:HG22	1:L:163:TRP:CZ3	2.48	0.49
2:H:136:VAL:HG23	2:H:185:SER:CA	2.42	0.48
1:L:54:ARG:HG3	1:L:54:ARG:HH11	1.78	0.48
1:L:163:TRP:N	1:L:163:TRP:CD1	2.81	0.48
1:L:108:ARG:CD	1:L:109:ALA:O	2.61	0.47
1:L:15:LEU:HD22	1:L:15:LEU:H	1.79	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:136:VAL:HG23	2:H:185:SER:OG	2.15	0.47
1:L:83:LEU:HD23	1:L:84:GLY:N	2.29	0.47
1:L:89:PHE:CZ	1:L:96:PHE:HB3	2.50	0.47
1:L:156:GLN:HE21	1:L:156:GLN:HB3	1.52	0.47
2:H:42:GLY:O	2:H:43:LYS:HG2	2.14	0.47
1:L:83:LEU:HD12	5:L:1028:HOH:O	2.15	0.46
2:H:171:GLN:HB2	2:H:173:ASP:OD1	2.15	0.45
2:H:24:ALA:HB1	2:H:27:TYR:CE1	2.52	0.45
3:A:4:DSN:N	3:A:8:DAS:OD2	2.50	0.44
1:L:49:TYR:CE2	4:L:1001:CIT:O6	2.70	0.43
1:L:94:ILE:HD13	1:L:96:PHE:CE1	2.54	0.43
2:H:46:LYS:HB3	2:H:46:LYS:HE3	1.68	0.42
1:L:119:PRO:HB3	1:L:209:PHE:CE2	2.54	0.42
3:A:1:DVA:HA	3:A:3:GLY:O	2.19	0.42
3:A:1:DVA:CA	3:A:3:GLY:H	2.32	0.42
1:L:54:ARG:HG3	1:L:54:ARG:NH1	2.35	0.42
1:L:119:PRO:HB3	1:L:209:PHE:CZ	2.55	0.41
1:L:204:PRO:HG2	5:L:1119:HOH:O	2.21	0.41
2:H:136:VAL:CG2	2:H:185:SER:HA	2.48	0.41
1:L:199:LYS:HB2	1:L:199:LYS:HE3	1.88	0.41
2:H:4:LEU:HD23	2:H:4:LEU:HA	1.89	0.41
3:A:8:DAS:O	3:A:9:DSN:HB3	2.21	0.41
3:A:9:DSN:HB3	5:A:143:HOH:O	2.21	0.41
1:L:27(B):ILE:HD11	1:L:71:PHE:CE2	2.56	0.41
1:L:35:TRP:CE2	1:L:73:LEU:HB2	2.56	0.41
2:H:173:ASP:C	2:H:174:LEU:HG	2.41	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	214/216 (99%)	210 (98%)	4 (2%)	0	100	100
2	H	205/216 (95%)	198 (97%)	7 (3%)	0	100	100
3	A	1/9 (11%)	1 (100%)	0	0	100	100
All	All	420/441 (95%)	409 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	194/194 (100%)	186 (96%)	8 (4%)	30	14
2	H	182/186 (98%)	177 (97%)	5 (3%)	44	28
All	All	376/380 (99%)	363 (96%)	13 (4%)	36	19

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	3	LEU
1	L	34	GLU
1	L	74	LYS
1	L	108	ARG
1	L	156	GLN
1	L	179	LEU
1	L	202	THR
1	L	210	ASN
2	H	80	LEU
2	H	149	PRO
2	H	150	VAL
2	H	174	LEU
2	H	177	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	156	GLN
2	H	3	GLN
2	H	81	GLN
2	H	82(A)	ASN
2	H	82(B)	ASN
2	H	171	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	CIT	L	1001	-	12,12,12	1.71	1 (8%)	17,17,17	3.91	8 (47%)
4	CIT	L	1002	-	12,12,12	1.50	1 (8%)	17,17,17	3.77	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	L	1001	-	-	9/16/16/16	-
4	CIT	L	1002	-	-	5/16/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	1001	CIT	C4-C3	3.73	1.58	1.53
4	L	1002	CIT	C4-C3	2.81	1.57	1.53

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	L	1001	CIT	C4-C3-C2	-8.25	87.64	109.16
4	L	1002	CIT	C4-C3-C2	-6.97	90.97	109.16
4	L	1002	CIT	O7-C3-C6	6.43	117.89	108.86
4	L	1001	CIT	C2-C3-C6	6.42	123.91	110.11
4	L	1002	CIT	O7-C3-C4	-6.26	94.74	109.40
4	L	1002	CIT	C2-C3-C6	5.97	122.94	110.11
4	L	1002	CIT	C4-C3-C6	-5.96	97.31	110.11
4	L	1001	CIT	O7-C3-C4	-5.94	95.49	109.40
4	L	1001	CIT	O7-C3-C6	5.71	116.89	108.86
4	L	1001	CIT	C4-C3-C6	-5.33	98.66	110.11
4	L	1001	CIT	O6-C6-C3	4.27	120.46	113.05
4	L	1002	CIT	O6-C6-C3	3.61	119.33	113.05
4	L	1001	CIT	O7-C3-C2	3.61	117.84	109.40
4	L	1002	CIT	O7-C3-C2	3.45	117.47	109.40
4	L	1001	CIT	O4-C5-C4	2.15	121.25	114.35

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	1001	CIT	C1-C2-C3-O7
4	L	1001	CIT	C1-C2-C3-C4
4	L	1001	CIT	C2-C3-C6-O5
4	L	1001	CIT	C2-C3-C6-O6
4	L	1001	CIT	O7-C3-C6-O5
4	L	1001	CIT	O7-C3-C6-O6
4	L	1002	CIT	C2-C3-C6-O5
4	L	1002	CIT	C2-C3-C6-O6
4	L	1001	CIT	C1-C2-C3-C6
4	L	1002	CIT	C1-C2-C3-O7
4	L	1002	CIT	C4-C3-C6-O6
4	L	1002	CIT	C4-C3-C6-O5
4	L	1001	CIT	C3-C4-C5-O4
4	L	1001	CIT	C3-C4-C5-O3

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	1001	CIT	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	216/216 (100%)	0.24	13 (6%) 21 21	13, 20, 38, 53	0
2	H	209/216 (96%)	-0.04	5 (2%) 59 58	10, 18, 31, 54	0
3	A	1/9 (11%)	1.55	0 100 100	27, 27, 27, 27	0
All	All	426/441 (96%)	0.10	18 (4%) 36 34	10, 19, 37, 54	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	41	GLY	6.1
2	H	173	ASP	4.1
2	H	43	LYS	3.6
2	H	174	LEU	3.6
2	H	42	GLY	3.5
1	L	9	LEU	3.2
1	L	169	LYS	2.8
1	L	202	THR	2.8
1	L	132	VAL	2.5
2	H	41	PRO	2.5
1	L	59	PRO	2.3
1	L	163	TRP	2.2
1	L	40	PRO	2.2
1	L	133	VAL	2.1
1	L	35	TRP	2.1
1	L	134	CYS	2.1
1	L	168	SER	2.1
1	L	73	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	DVA	A	1	7/8	0.37	0.38	27,49,60,63	0
3	DSN	A	9	7/7	0.68	0.43	53,60,64,66	0
3	DAS	A	8	8/9	0.76	0.18	30,46,49,59	0
3	DPR	A	2	7/8	0.82	0.33	40,51,53,56	0
3	DTY	A	7	12/13	0.93	0.11	17,18,23,27	0
3	DHI	A	6	10/11	0.95	0.09	14,15,18,19	0
3	DSN	A	4	6/7	0.96	0.06	13,18,21,25	0
3	DGN	A	5	9/10	0.97	0.07	10,15,22,22	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CIT	L	1002	13/13	0.46	0.40	72,83,89,91	0
4	CIT	L	1001	13/13	0.73	0.40	35,60,82,85	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.