



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 03:12 pm GMT

PDB ID : 4ZH5  
Title : Crystal structure of Endoglucanase from *Perinereis brevicirris* with Cellobiose  
Authors : Fewings, R.S.; Swiderska, A.; Sanchez-Weatherby, J.; Sorensen, T.-M.; Schnorr, K.M.; Kneale, G.G.; McGeehan, J.E.  
Deposited on : 2015-04-24  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	426	 93% 6%
1	B	426	 94% 6%
2	C	2	 100%
2	D	2	 50% 50%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7961 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoglucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	425	3407	2139	581	672	15	0	16	0
1	B	425	3382	2124	576	668	14	0	13	0

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	23	12	11	0	0	0
2	D	2	23	12	11	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0
3	B	1	1	1	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	A	4	4	4	0	0
4	B	1	1	1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Cl 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	593	Total O 593 593	0	0
6	B	525	Total O 525 525	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endoglucanase

Chain A:  93% 6%




- Molecule 1: Endoglucanase

Chain B:  94% 6%



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain C:  100%



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain D:  50% 50%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.15Å 108.17Å 112.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	16.98 – 1.35 16.98 – 1.35	Depositor EDS
% Data completeness (in resolution range)	97.7 (16.98-1.35) 97.8 (16.98-1.35)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 1.35Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.115 , 0.148 0.115 , 0.149	Depositor DCC
$R_{free}$ test set	9867 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	9.3	Xtrriage
Anisotropy	0.529	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.017 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	7961	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, CA, CL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	1/3520 (0.0%)	0.90	14/4800 (0.3%)
1	B	0.72	2/3498 (0.1%)	0.91	13/4771 (0.3%)
All	All	0.74	3/7018 (0.0%)	0.91	27/9571 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	414	GLU	CD-OE2	5.86	1.32	1.25
1	B	217	GLU	CD-OE1	5.73	1.31	1.25
1	B	414	GLU	CD-OE2	5.47	1.31	1.25

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	102	ARG	NE-CZ-NH2	-10.46	115.07	120.30
1	A	102	ARG	NE-CZ-NH2	-10.35	115.12	120.30
1	B	102	ARG	NE-CZ-NH1	8.81	124.71	120.30
1	A	102	ARG	NE-CZ-NH1	8.26	124.43	120.30
1	A	94	ARG	NE-CZ-NH2	-7.92	116.34	120.30
1	A	274	ARG	NE-CZ-NH2	-7.38	116.61	120.30
1	B	94	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	A	419	TYR	CB-CG-CD1	6.22	124.73	121.00
1	B	94	ARG	NE-CZ-NH2	-6.21	117.19	120.30
1	B	115	ASP	CB-CG-OD1	6.02	123.72	118.30
1	A	274	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	B	54	ASP	CB-CG-OD1	5.91	123.61	118.30
1	A	285	PHE	CB-CG-CD1	5.77	124.84	120.80
1	A	197	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	B	359	ARG	NE-CZ-NH2	5.68	123.14	120.30
1	B	419	TYR	CB-CG-CD1	5.57	124.34	121.00

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	334	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	B	419	TYR	CB-CG-CD2	-5.43	117.74	121.00
1	B	341	LEU	CB-CG-CD1	5.36	120.11	111.00
1	A	124	ASP	CB-CG-OD1	5.20	122.98	118.30
1	B	10	TYR	CB-CG-CD2	5.16	124.10	121.00
1	B	13	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	A	94	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	B	38	ASP	CB-CG-OD1	5.08	122.87	118.30
1	A	138	MET	CG-SD-CE	5.07	108.31	100.20
1	A	197	ARG	NE-CZ-NH1	5.07	122.83	120.30
1	A	414	GLU	N-CA-CB	5.03	119.65	110.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3407	0	3116	37	0
1	B	3382	0	3092	30	0
2	C	23	0	21	0	0
2	D	23	0	21	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	0	0
4	B	1	0	0	0	0
5	B	1	0	0	0	0
6	A	593	0	0	10	0
6	B	525	0	0	11	0
All	All	7961	0	6250	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:337[B]:LEU:CD1	1:A:430[B]:ILE:HD12	1.30	1.60
1:A:337[B]:LEU:HD12	1:A:430[B]:ILE:CD1	1.09	1.54
1:A:337[B]:LEU:CD1	1:A:430[B]:ILE:CD1	1.80	1.53
1:A:303[A]:TRP:CZ2	1:B:303[A]:TRP:CZ2	2.10	1.37
1:A:303[A]:TRP:HZ2	1:B:303[A]:TRP:CE2	1.48	1.28
1:A:303[A]:TRP:CZ2	1:B:303[A]:TRP:CE2	2.22	1.27
1:A:303[A]:TRP:CE2	1:B:303[A]:TRP:HZ2	1.55	1.23
1:A:303[A]:TRP:CE2	1:B:303[A]:TRP:CZ2	2.27	1.19
1:A:303[A]:TRP:CZ2	1:B:303[A]:TRP:NE1	2.24	1.05
1:A:337[B]:LEU:HD13	1:A:430[B]:ILE:HD12	1.06	1.05
1:A:303[A]:TRP:NE1	1:B:303[A]:TRP:CZ2	2.29	1.00
1:A:337[B]:LEU:CD1	1:A:430[B]:ILE:HD11	1.66	0.96
1:B:321[A]:GLU:OE2	1:B:330:ARG:NH2	2.07	0.86
1:A:412[A]:SER:OG	6:A:601:HOH:O	1.98	0.81
1:B:9:GLN:N	6:B:602:HOH:O	2.19	0.76
1:A:411:ILE:O	6:A:603:HOH:O	2.04	0.75
1:A:217[A]:GLU:OE1	6:A:602:HOH:O	2.03	0.75
1:A:303[A]:TRP:CE2	1:B:303[A]:TRP:CE2	2.66	0.75
1:A:303[A]:TRP:CE2	1:B:303[A]:TRP:NE1	2.58	0.72
1:A:337[B]:LEU:HD12	1:A:430[B]:ILE:HD11	0.69	0.68
1:A:303[B]:TRP:CD1	6:A:603:HOH:O	2.45	0.68
1:B:303[B]:TRP:CD1	6:B:604:HOH:O	2.47	0.68
1:A:303[A]:TRP:NE1	1:B:303[A]:TRP:CE2	2.61	0.68
1:A:303[A]:TRP:NE1	1:B:303[A]:TRP:HZ2	1.77	0.67
1:A:303[A]:TRP:CZ3	1:A:411:ILE:HG12	2.29	0.67
1:A:303[A]:TRP:HZ2	1:B:303[A]:TRP:CZ2	1.74	0.67
1:A:303[A]:TRP:HZ2	1:B:303[A]:TRP:NE1	1.75	0.65
1:B:412[A]:SER:OG	6:B:601:HOH:O	2.09	0.62
1:B:239:LYS:HE2	6:B:631:HOH:O	2.01	0.60
1:A:303[A]:TRP:CZ2	1:B:303[A]:TRP:HZ2	1.78	0.57
1:A:217[A]:GLU:OE2	6:A:604:HOH:O	2.18	0.55
1:A:409:ASP:CG	1:A:412[A]:SER:HB2	2.29	0.53
1:B:303[A]:TRP:CZ3	1:B:411:ILE:HG12	2.44	0.53
1:B:303[B]:TRP:HD1	6:B:604:HOH:O	1.90	0.52
1:B:181:GLN:HG3	6:B:942:HOH:O	2.10	0.51
1:A:303[A]:TRP:NE1	1:B:303[A]:TRP:NE1	2.61	0.49
1:A:337[B]:LEU:HD13	1:A:430[B]:ILE:CD1	1.91	0.48
1:A:217[B]:GLU:OE1	6:A:605:HOH:O	2.20	0.47
1:A:239:LYS:HE2	6:A:626:HOH:O	2.14	0.47
1:A:110[A]:LYS:NZ	6:A:607:HOH:O	2.38	0.47
1:B:321[B]:GLU:HG2	6:B:835:HOH:O	2.16	0.46
1:A:65:HIS:HE1	6:A:868:HOH:O	1.99	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:65:HIS:HD2	6:B:874:HOH:O	2.00	0.45
1:B:181:GLN:CG	6:B:942:HOH:O	2.64	0.45
1:A:303[B]:TRP:HD1	6:A:603:HOH:O	1.92	0.45
1:B:65:HIS:HE1	6:B:849:HOH:O	2.00	0.44
1:B:276[B]:GLN:NE2	6:B:616:HOH:O	2.50	0.44
1:A:411:ILE:CD1	1:B:411:ILE:HD11	2.48	0.42
1:A:314:PHE:CZ	1:A:318:LEU:HD11	2.54	0.41
1:A:411:ILE:HD11	1:B:411:ILE:CD1	2.50	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	439/426 (103%)	429 (98%)	9 (2%)	1 (0%)	47 21
1	B	436/426 (102%)	426 (98%)	9 (2%)	1 (0%)	47 21
All	All	875/852 (103%)	855 (98%)	18 (2%)	2 (0%)	47 21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	342	GLY
1	B	342	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	347/331 (105%)	341 (98%)	6 (2%)	60	28
1	B	344/331 (104%)	341 (99%)	3 (1%)	78	53
All	All	691/662 (104%)	682 (99%)	9 (1%)	71	37

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	155	ASP
1	A	346	ARG
1	A	412[A]	SER
1	A	412[B]	SER
1	A	413	ASN
1	A	418	ASP
1	B	155	ASP
1	B	346	ARG
1	B	413	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	65	HIS
1	A	116	ASN
1	B	65	HIS
1	B	116	ASN
1	B	180	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BGC	C	1	2	12,12,12	0.73	0	17,17,17	1.47	2 (11%)
2	BGC	C	2	2	11,11,12	0.88	0	15,15,17	1.43	2 (13%)
2	BGC	D	1	2	12,12,12	0.78	0	17,17,17	0.87	0
2	BGC	D	2	2	11,11,12	0.39	0	15,15,17	1.34	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	0/2/22/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	BGC	D	1	2	-	0/2/22/22	0/1/1/1
2	BGC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	BGC	C4-C3-C2	-4.25	103.41	110.82
2	C	2	BGC	O3-C3-C2	-3.97	102.40	109.99
2	D	2	BGC	O3-C3-C2	-3.22	103.83	109.99
2	C	2	BGC	O3-C3-C4	2.42	115.94	110.35
2	C	1	BGC	O5-C1-C2	-2.21	106.34	110.28


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	425/426 (99%)	-0.50	1 (0%) 95 95	5, 9, 17, 57	0
1	B	425/426 (99%)	-0.34	2 (0%) 91 92	6, 13, 25, 34	0
All	All	850/852 (99%)	-0.42	3 (0%) 92 93	5, 11, 23, 57	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	410	TYR	4.2
1	B	371	PRO	2.7
1	A	9	GLN	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	C	2	11/12	0.92	0.10	14,17,23,23	11
2	BGC	C	1	12/12	0.94	0.09	15,17,22,24	12
2	BGC	D	1	12/12	0.95	0.09	11,15,20,22	12
2	BGC	D	2	11/12	0.96	0.07	11,13,16,18	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NA	B	505	1/1	0.92	0.09	41,41,41,41	0
4	NA	A	505	1/1	0.99	0.22	29,29,29,29	0
4	NA	A	507	1/1	0.99	0.14	33,33,33,33	0
4	NA	A	504	1/1	0.99	0.13	34,34,34,34	0
4	NA	A	506	1/1	1.00	0.03	16,16,16,16	0
3	CA	A	501	1/1	1.00	0.04	7,7,7,7	0
3	CA	B	503	1/1	1.00	0.04	6,6,6,6	0
5	CL	B	504	1/1	1.00	0.03	16,16,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.