



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 12, 2023 – 02:25 pm GMT

PDB ID : 3ZKX
Title : TERNARY BACE2 XAPERONE COMPLEX
Authors : Kuglstatter, A.; Banner, D.W.; Benz, J.; Bertschinger, J.; Burger, D.; Cuppuleri, S.; Debulpaep, M.; Gast, A.; Grabulovski, D.; Gsell, B.; Hilpert, H.; Huber, W.; Kuszniir, E.; Laeremans, T.; Matile, H.; Rufer, A.; Schlatter, D.; Steyeart, J.; Stihle, M.; Thoma, R.; Weber, M.; Ruf, A.
Deposited on : 2013-01-25
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 4%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 30px;">3% 88% 7% •</p>
2	B	122	<div style="display: flex; align-items: center;"> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 30px;">84% 6% • 9%</p>
3	C	122	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 73%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 30px;">3% 73% 16% •• 9%</p>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-SECRETASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	370	2866	1848	456	550	12	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	269	ALA	GLU	engineered mutation	UNP Q9Y5Z0

- Molecule 2 is a protein called XA4813.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	111	827	509	150	164	4	0	0	0

- Molecule 3 is a protein called XA4815.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	111	848	521	165	158	4	0	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
5	B	1	4	2	1	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	202	Total	O	0	0
			202	202		
6	B	53	Total	O	0	0
			53	53		
6	C	56	Total	O	0	0
			56	56		

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	63.37Å 153.83Å 247.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.70 – 2.37 48.17 – 2.37	Depositor EDS
% Data completeness (in resolution range)	99.6 (34.70-2.37) 99.6 (48.17-2.37)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.37Å)	Xtrriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.185 , 0.208 0.182 , 0.210	Depositor DCC
R_{free} test set	2504 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	56.6	Xtrriage
Anisotropy	0.342	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 61.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4857	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2937	0.70	0/3996
2	B	0.50	0/841	0.73	0/1138
3	C	0.52	0/865	0.77	0/1166
All	All	0.51	0/4643	0.72	0/6300

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2866	0	2789	12	0
2	B	827	0	809	5	0
3	C	848	0	809	9	0
4	B	1	0	0	0	0
5	B	4	0	6	0	0
6	A	202	0	0	0	0
6	B	53	0	0	0	0
6	C	56	0	0	0	0
All	All	4857	0	4413	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:193:MET:HG2	3:C:238:LEU:HD22	1.77	0.66
1:A:295:LYS:HG2	1:A:314:LEU:HD22	1.92	0.52
1:A:283:TRP:CZ3	1:A:289:PRO:HB3	2.47	0.49
3:C:191:ALA:HB3	3:C:257:TYR:CE1	2.48	0.49
3:C:256:ILE:HG12	3:C:261:ARG:HG3	1.94	0.49
2:B:210:ILE:HG13	2:B:217:THR:HG22	1.95	0.49
3:C:191:ALA:HB1	3:C:258:ASN:HB2	1.94	0.49
2:B:242:MET:HE2	2:B:245:LEU:HD21	1.95	0.48
1:A:36:ILE:HD12	1:A:43:LEU:HD13	1.94	0.48
1:A:283:TRP:CH2	1:A:289:PRO:HB3	2.49	0.48
3:C:238:LEU:HD11	3:C:257:TYR:CD1	2.49	0.47
2:B:254:TYR:CD2	2:B:261:ARG:HD2	2.51	0.46
1:A:316:GLN:HB3	1:A:374:ILE:HD12	1.99	0.45
1:A:126:LEU:O	1:A:129:ILE:HG12	2.16	0.45
2:B:242:MET:HB3	2:B:245:LEU:HD21	2.00	0.44
1:A:61:HIS:HB3	1:A:64:ILE:HG13	2.00	0.43
1:A:34:MET:HE3	1:A:135:LEU:HD22	2.01	0.43
3:C:210:ILE:HD11	3:C:229:ALA:HB1	2.01	0.43
1:A:246:LEU:HD22	1:A:335:GLY:HA2	2.00	0.42
1:A:34:MET:CE	1:A:135:LEU:HD22	2.49	0.42
3:C:194:ARG:HG3	3:C:209:ASN:OD1	2.20	0.42
3:C:181:CYS:HB3	3:C:238:LEU:HB2	2.02	0.42
1:A:320:GLN:HA	1:A:321:PRO:HD3	1.96	0.42
1:A:264:ALA:HB3	1:A:294:PRO:HG3	2.02	0.41
2:B:160:GLN:HE21	2:B:160:GLN:HB3	1.75	0.41
3:C:174:GLY:HA2	3:C:244:ASN:HD22	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/386 (94%)	356 (98%)	8 (2%)	0	100	100
2	B	109/122 (89%)	107 (98%)	2 (2%)	0	100	100
3	C	109/122 (89%)	105 (96%)	3 (3%)	1 (1%)	17	23
All	All	582/630 (92%)	568 (98%)	13 (2%)	1 (0%)	47	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	258	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/319 (98%)	301 (97%)	10 (3%)	39	56
2	B	89/99 (90%)	87 (98%)	2 (2%)	52	69
3	C	84/94 (89%)	73 (87%)	11 (13%)	4	4
All	All	484/512 (94%)	461 (95%)	23 (5%)	25	39

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	175	LEU
1	A	198	LYS
1	A	210	TRP
1	A	237	LYS
1	A	320	GLN
1	A	333	ARG
1	A	341	ASN
1	A	371	CYS
1	A	391	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	160	GLN
2	B	204	ARG
3	C	163	LEU
3	C	171	VAL
3	C	172	GLN
3	C	204	LEU
3	C	223	VAL
3	C	231	ARG
3	C	233	LYS
3	C	238	LEU
3	C	246	ARG
3	C	258	ASN
3	C	261	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	341	ASN
3	C	244	ASN
3	C	258	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DMS	B	1272	-	3,3,3	0.37	0	3,3,3	0.67	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/386 (95%)	-0.06	13 (3%) 44 47	44, 62, 93, 133	0
2	B	111/122 (90%)	-0.23	0 100 100	50, 64, 89, 108	0
3	C	111/122 (90%)	0.19	4 (3%) 42 46	44, 64, 96, 126	0
All	All	592/630 (93%)	-0.04	17 (2%) 51 53	44, 63, 93, 133	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	272	HIS	4.8
3	C	273	HIS	4.5
1	A	16	LEU	4.1
1	A	235	ALA	3.3
3	C	214	ASP	3.0
1	A	227	ASN	3.0
1	A	108	ASN	2.8
1	A	341	ASN	2.8
1	A	107	PHE	2.6
1	A	119	PHE	2.6
1	A	127	PRO	2.5
1	A	63	TYR	2.5
1	A	175	LEU	2.4
3	C	271	SER	2.4
1	A	236	ASP	2.2
1	A	128	GLY	2.2
1	A	104	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CL	B	1271	1/1	0.89	0.26	82,82,82,82	0
5	DMS	B	1272	4/4	0.95	0.17	65,71,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.