



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2023 – 02:24 AM EDT

PDB ID : 2ZO9  
Title : Malonate-bound structure of the glycerophosphodiesterase from *Enterobacter aerogenes* (GpdQ) and characterization of the native Fe<sup>2+</sup> metal ion preference  
Authors : Jackson, C.J.; Carr, P.D.; Ollis, D.L.  
Deposited on : 2008-05-07  
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	274	 11% 80% 14% . .
1	C	274	 10% 80% 14% . .

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	271	2133	1348	372	399	14	0	0	0
1	C	271	2133	1348	372	399	14	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	224	ALA	GLU	SEE REMARK 999	UNP Q6XBH1
B	227	ASP	ARG	SEE REMARK 999	UNP Q6XBH1
C	224	ALA	GLU	SEE REMARK 999	UNP Q6XBH1
C	227	ASP	ARG	SEE REMARK 999	UNP Q6XBH1

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Fe	0	0
			2	2		
2	C	2	Total	Fe	0	0
			2	2		

- Molecule 3 is MALONATE ION (three-letter code: MLI) (formula: C<sub>3</sub>H<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 7 3 4	0	0
3	C	1	Total C O 7 3 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	80	Total O 80 80	0	0
4	C	81	Total O 81 81	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphohydrolase



- Molecule 1: Phosphohydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	164.16Å 164.16Å 164.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.03 – 2.20 29.02 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.2 (29.03-2.20) 98.2 (29.02-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.43 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.170 , 0.189 0.169 , 0.187	Depositor DCC
$R_{free}$ test set	3703 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.8	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 48.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.031 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4445	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.69	0/2192	0.83	8/2991 (0.3%)
1	C	0.71	0/2192	0.82	8/2991 (0.3%)
All	All	0.70	0/4384	0.83	16/5982 (0.3%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	190	ARG	NE-CZ-NH2	-10.44	115.08	120.30
1	B	208	LEU	CA-CB-CG	7.75	133.13	115.30
1	C	208	LEU	CA-CB-CG	7.62	132.84	115.30
1	B	12	ARG	NE-CZ-NH2	-7.40	116.60	120.30
1	C	12	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	B	190	ARG	NE-CZ-NH1	6.74	123.67	120.30
1	B	190	ARG	CG-CD-NE	-6.56	98.03	111.80
1	C	190	ARG	NE-CZ-NH2	-6.27	117.17	120.30
1	B	106	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	B	12	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	C	12	ARG	NE-CZ-NH1	5.17	122.88	120.30
1	B	139	LEU	CA-CB-CG	5.10	127.03	115.30
1	C	30	ASP	CB-CG-OD1	5.07	122.87	118.30
1	C	190	ARG	NE-CZ-NH1	5.05	122.83	120.30
1	C	3	LEU	CA-CB-CG	5.04	126.89	115.30
1	C	36	ASN	N-CA-CB	5.02	119.64	110.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2133	0	2046	54	0
1	C	2133	0	2046	45	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	B	7	0	2	0	0
3	C	7	0	2	0	0
4	B	80	0	0	3	0
4	C	81	0	0	5	0
All	All	4445	0	4096	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:152:THR:HB	1:B:190:ARG:HB2	1.39	0.98
1:C:152:THR:HB	1:C:190:ARG:HB2	1.46	0.97
1:B:166:GLN:HG2	1:B:167:MET:HE3	1.48	0.95
1:B:131:LEU:H	1:B:174:ASN:HD21	1.17	0.89
1:C:131:LEU:H	1:C:174:ASN:HD21	1.13	0.89
1:B:190:ARG:NH2	1:C:203:GLN:HB3	1.89	0.88
1:B:203:GLN:HG3	1:C:208:LEU:HB2	1.58	0.85
1:B:5:HIS:HD2	1:B:240:HIS:HE1	1.25	0.82
1:C:5:HIS:HD2	1:C:240:HIS:HE1	1.27	0.79
1:B:203:GLN:HB2	1:C:190:ARG:NH2	2.00	0.77
1:C:5:HIS:CD2	1:C:240:HIS:HE1	2.03	0.76
1:B:5:HIS:CD2	1:B:240:HIS:HE1	2.04	0.76
1:B:39:ARG:HH11	1:B:39:ARG:HG2	1.52	0.74
1:B:166:GLN:HG2	1:B:167:MET:CE	2.20	0.71
1:B:110:ASP:HB2	1:B:149:LYS:NZ	2.05	0.70
1:C:197:HIS:HB2	1:C:219:VAL:HG13	1.73	0.69
1:C:200:THR:HG21	4:C:299:HOH:O	1.94	0.66
1:B:174:ASN:HD22	1:B:174:ASN:H	1.44	0.66
1:B:166:GLN:CG	1:B:167:MET:HE3	2.26	0.65

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:ASP:HA	4:B:285:HOH:O	1.96	0.65
1:C:36:ASN:ND2	1:C:71:ASN:H	1.94	0.65
1:C:190:ARG:NH1	1:C:247:TRP:HB3	2.12	0.65
1:B:200:THR:HG21	4:B:284:HOH:O	1.99	0.63
1:B:203:GLN:HE22	1:C:203:GLN:CG	2.10	0.62
1:B:110:ASP:HB2	1:B:149:LYS:HZ3	1.64	0.61
1:C:7:SER:HB3	1:C:48:SER:OG	2.01	0.61
1:C:252:HIS:HE1	4:C:322:HOH:O	1.83	0.60
1:B:250:TYR:HB3	1:C:200:THR:HB	1.84	0.60
1:C:240:HIS:HD2	1:C:249:SER:OG	1.85	0.59
1:B:97:GLN:HE21	1:B:97:GLN:H	1.51	0.57
1:B:197:HIS:HB2	1:B:219:VAL:HG13	1.86	0.57
1:B:5:HIS:HD2	1:B:240:HIS:CE1	2.16	0.57
1:B:152:THR:HB	1:B:190:ARG:CB	2.26	0.57
1:B:203:GLN:HB2	1:C:190:ARG:HH21	1.69	0.57
1:B:18:LEU:HD22	1:B:19:TYR:CE2	2.42	0.55
1:B:36:ASN:HD21	1:B:71:ASN:H	1.54	0.55
1:B:36:ASN:ND2	1:B:71:ASN:H	2.03	0.55
1:B:203:GLN:NE2	1:C:203:GLN:HB2	2.21	0.55
1:C:36:ASN:HD21	1:C:70:LEU:HA	1.72	0.55
1:C:36:ASN:HB3	4:C:316:HOH:O	2.06	0.55
1:B:252:HIS:HE1	4:B:286:HOH:O	1.89	0.55
1:C:5:HIS:HD2	1:C:240:HIS:CE1	2.17	0.55
1:B:164:ASN:HD22	1:B:164:ASN:C	2.10	0.54
1:B:200:THR:HB	1:C:250:TYR:HB3	1.90	0.53
1:B:39:ARG:HG2	1:B:39:ARG:NH1	2.22	0.53
1:B:106:ARG:NH2	1:B:134:GLU:OE2	2.41	0.53
1:C:152:THR:HG21	1:C:247:TRP:CZ3	2.44	0.53
1:C:5:HIS:HE1	1:C:48:SER:OG	1.91	0.52
1:C:36:ASN:HD21	1:C:71:ASN:H	1.56	0.51
1:B:190:ARG:NH2	1:C:203:GLN:OE1	2.44	0.50
1:C:123:ARG:CG	1:C:123:ARG:HH11	2.25	0.50
1:B:203:GLN:HE22	1:C:203:GLN:CB	2.25	0.50
1:B:206:GLN:NE2	1:B:206:GLN:H	2.09	0.49
1:B:263:TYR:HB3	1:C:228:PRO:HB2	1.94	0.49
1:C:104:ASN:ND2	1:C:106:ARG:HH22	2.11	0.49
1:C:174:ASN:HD22	1:C:174:ASN:H	1.59	0.49
1:B:190:ARG:HH22	1:C:203:GLN:HB3	1.73	0.48
1:B:152:THR:HG21	1:B:247:TRP:CZ3	2.48	0.48
1:B:240:HIS:HD2	1:B:249:SER:OG	1.96	0.48
1:C:155:MET:O	1:C:193:CYS:HA	2.14	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:ARG:NH2	1:B:134:GLU:HG2	2.29	0.47
1:B:208:LEU:HD22	1:C:208:LEU:HD22	1.96	0.46
1:C:152:THR:CG2	1:C:247:TRP:CZ3	2.98	0.46
1:B:230:TYR:CZ	1:C:261:TRP:HB2	2.51	0.46
1:B:104:ASN:HD22	1:B:106:ARG:HH12	1.63	0.46
1:B:203:GLN:HG2	1:B:208:LEU:HA	1.96	0.46
1:B:5:HIS:HE1	1:B:48:SER:OG	1.99	0.46
1:B:49:GLY:HA2	1:B:77:ILE:O	2.16	0.45
1:B:155:MET:O	1:B:193:CYS:HA	2.17	0.45
1:C:198:SER:OG	1:C:200:THR:HG22	2.16	0.44
1:B:138:TRP:O	1:B:142:GLN:HG2	2.16	0.44
1:C:18:LEU:HD22	1:C:19:TYR:CE2	2.54	0.43
1:C:164:ASN:C	1:C:164:ASN:HD22	2.22	0.42
1:C:240:HIS:CD2	1:C:249:SER:OG	2.68	0.42
1:B:261:TRP:HB2	1:C:230:TYR:CZ	2.55	0.42
1:B:203:GLN:HE22	1:C:203:GLN:HG2	1.84	0.41
1:B:18:LEU:HD22	1:B:19:TYR:CD2	2.56	0.41
1:B:48:SER:HA	1:B:77:ILE:HG12	2.02	0.41
1:B:240:HIS:CD2	1:B:247:TRP:HE1	2.39	0.41
1:C:265:GLU:HB2	4:C:358:HOH:O	2.21	0.41
1:B:198:SER:OG	1:B:200:THR:HG22	2.22	0.40
1:C:206:GLN:H	1:C:206:GLN:NE2	2.20	0.40
1:C:152:THR:HG22	4:C:289:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	269/274 (98%)	261 (97%)	7 (3%)	1 (0%)	34	37
1	C	269/274 (98%)	260 (97%)	8 (3%)	1 (0%)	34	37

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	538/548 (98%)	521 (97%)	15 (3%)	2 (0%)	34 37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	216	VAL
1	C	216	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	232/235 (99%)	214 (92%)	18 (8%)	12 13
1	C	232/235 (99%)	208 (90%)	24 (10%)	7 6
All	All	464/470 (99%)	422 (91%)	42 (9%)	9 9

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	3	LEU
1	B	12	ARG
1	B	18	LEU
1	B	39	ARG
1	B	58	GLU
1	B	97	GLN
1	B	123	ARG
1	B	139	LEU
1	B	152	THR
1	B	164	ASN
1	B	174	ASN
1	B	179	LEU
1	B	181	LEU
1	B	182	VAL
1	B	200	THR
1	B	206	GLN

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	B	208	LEU
1	B	238	LEU
1	C	3	LEU
1	C	12	ARG
1	C	14	ARG
1	C	18	LEU
1	C	36	ASN
1	C	84	LYS
1	C	115	ARG
1	C	123	ARG
1	C	139	LEU
1	C	145	GLU
1	C	152	THR
1	C	160	LEU
1	C	164	ASN
1	C	174	ASN
1	C	179	LEU
1	C	182	VAL
1	C	184	ARG
1	C	187	SER
1	C	200	THR
1	C	205	ARG
1	C	206	GLN
1	C	208	LEU
1	C	226	THR
1	C	238	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	B	5	HIS
1	B	28	ASN
1	B	34	GLN
1	B	36	ASN
1	B	97	GLN
1	B	104	ASN
1	B	164	ASN
1	B	174	ASN
1	B	203	GLN
1	B	206	GLN
1	B	223	HIS
1	B	240	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	242	GLN
1	B	251	GLN
1	C	5	HIS
1	C	28	ASN
1	C	36	ASN
1	C	61	GLN
1	C	97	GLN
1	C	104	ASN
1	C	142	GLN
1	C	164	ASN
1	C	166	GLN
1	C	174	ASN
1	C	206	GLN
1	C	223	HIS
1	C	240	HIS
1	C	242	GLN
1	C	251	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MLI	C	277	-	6,6,6	1.07	0	7,7,7	1.14	0
3	MLI	B	277	-	6,6,6	1.13	0	7,7,7	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	C	277	-	-	3/4/4/4	-
3	MLI	B	277	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	277	MLI	C3-C1-C2-O6
3	B	277	MLI	C3-C1-C2-O6
3	C	277	MLI	C3-C1-C2-O7
3	B	277	MLI	C3-C1-C2-O7
3	C	277	MLI	C2-C1-C3-O8

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	271/274 (98%)	0.20	29 (10%) <b>6</b>   <b>5</b>	43, 50, 58, 67	0
1	C	271/274 (98%)	0.22	28 (10%) <b>6</b>   <b>5</b>	42, 50, 58, 65	0
All	All	542/548 (98%)	0.21	57 (10%) <b>6</b>   <b>5</b>	42, 50, 58, 67	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	144	PHE	5.5
1	B	148	ASP	5.2
1	C	148	ASP	4.6
1	C	224	ALA	4.1
1	B	47	VAL	3.7
1	C	125	GLY	3.6
1	C	48	SER	3.5
1	C	271	THR	3.5
1	C	145	GLU	3.5
1	B	77	ILE	3.4
1	C	47	VAL	3.4
1	B	49	GLY	3.4
1	C	154	PHE	3.4
1	C	194	GLY	3.3
1	B	125	GLY	3.2
1	C	124	ALA	3.1
1	B	192	PHE	3.1
1	C	77	ILE	3.0
1	B	6	ILE	3.0
1	B	51	ILE	3.0
1	B	224	ALA	3.0
1	B	226	THR	3.0
1	B	153	ILE	3.0
1	C	153	ILE	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	100	SER	3.0
1	C	193	CYS	3.0
1	B	48	SER	2.9
1	B	212	LEU	2.9
1	B	46	VAL	2.9
1	C	100	SER	2.9
1	C	155	MET	2.9
1	B	191	ILE	2.8
1	C	119	ILE	2.8
1	B	110	ASP	2.8
1	C	191	ILE	2.8
1	B	154	PHE	2.8
1	B	119	ILE	2.8
1	C	192	PHE	2.7
1	B	155	MET	2.6
1	B	245	GLU	2.6
1	C	6	ILE	2.6
1	B	271	THR	2.6
1	C	147	GLY	2.6
1	C	7	SER	2.4
1	C	226	THR	2.4
1	C	223	HIS	2.3
1	C	49	GLY	2.3
1	C	209	ILE	2.2
1	C	211	THR	2.2
1	B	147	GLY	2.2
1	B	99	GLY	2.2
1	B	7	SER	2.2
1	B	124	ALA	2.1
1	B	225	ASP	2.1
1	C	238	LEU	2.1
1	B	193	CYS	2.1
1	B	238	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MLI	B	277	7/7	0.90	0.30	64,65,67,67	7
3	MLI	C	277	7/7	0.92	0.33	61,62,63,63	7
2	FE2	C	275	1/1	0.99	0.03	45,45,45,45	1
2	FE2	B	275	1/1	0.99	0.02	44,44,44,44	1
2	FE2	B	276	1/1	0.99	0.02	47,47,47,47	1
2	FE2	C	276	1/1	1.00	0.05	47,47,47,47	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.