



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2024 – 02:57 PM JST

PDB ID : 4ZRC  
Title : Crystal structure of MSM-13, a putative T1-like thiolase from *Mycobacterium smegmatis*  
Authors : Janardan, N.; Harijan, R.K.; Keima, T.R.; Wierenga, R.; Murthy, M.R.N.  
Deposited on : 2015-05-12  
Resolution : 2.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	 2% 77% 15% .. 5%
1	B	413	 2% 77% 16% .. .
1	C	413	 2% 76% 18% . .
1	D	413	 3% 74% 19% . .

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 11586 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-ketothiolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	391	Total 2794	C 1721	N 513	O 544	S 16	0	0	0
1	B	397	Total 2885	C 1780	N 531	O 558	S 16	0	0	0
1	C	396	Total 2904	C 1790	N 537	O 561	S 16	0	0	0
1	D	395	Total 2900	C 1790	N 535	O 559	S 16	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP A0QUH3
A	-4	HIS	-	expression tag	UNP A0QUH3
A	-3	HIS	-	expression tag	UNP A0QUH3
A	-2	HIS	-	expression tag	UNP A0QUH3
A	-1	HIS	-	expression tag	UNP A0QUH3
A	0	HIS	-	expression tag	UNP A0QUH3
B	-5	HIS	-	expression tag	UNP A0QUH3
B	-4	HIS	-	expression tag	UNP A0QUH3
B	-3	HIS	-	expression tag	UNP A0QUH3
B	-2	HIS	-	expression tag	UNP A0QUH3
B	-1	HIS	-	expression tag	UNP A0QUH3
B	0	HIS	-	expression tag	UNP A0QUH3
C	-5	HIS	-	expression tag	UNP A0QUH3
C	-4	HIS	-	expression tag	UNP A0QUH3
C	-3	HIS	-	expression tag	UNP A0QUH3
C	-2	HIS	-	expression tag	UNP A0QUH3
C	-1	HIS	-	expression tag	UNP A0QUH3
C	0	HIS	-	expression tag	UNP A0QUH3
D	-5	HIS	-	expression tag	UNP A0QUH3
D	-4	HIS	-	expression tag	UNP A0QUH3
D	-3	HIS	-	expression tag	UNP A0QUH3

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	HIS	-	expression tag	UNP A0QUH3
D	-1	HIS	-	expression tag	UNP A0QUH3
D	0	HIS	-	expression tag	UNP A0QUH3





- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	31	Total O 31 31	0	0
2	B	25	Total O 25 25	0	0
2	C	32	Total O 32 32	0	0
2	D	15	Total O 15 15	0	0





- Molecule 1: Beta-ketothiolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.46Å 104.14Å 104.57Å 90.00° 107.53° 90.00°	Depositor
Resolution (Å)	47.52 – 2.70 69.05 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (47.52-2.70) 99.0 (69.05-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.07 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.223 , 0.283 0.223 , 0.283	Depositor DCC
$R_{free}$ test set	2129 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.1	Xtrriage
Anisotropy	0.066	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 17.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	11586	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.68	5/2831 (0.2%)	0.83	6/3837 (0.2%)
1	B	0.65	5/2927 (0.2%)	0.81	4/3970 (0.1%)
1	C	0.64	2/2944 (0.1%)	0.79	1/3987 (0.0%)
1	D	0.68	4/2942 (0.1%)	0.87	3/3989 (0.1%)
All	All	0.66	16/11644 (0.1%)	0.82	14/15783 (0.1%)

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	178	ARG	CZ-NH1	-12.12	1.17	1.33
1	D	126	TYR	CE1-CZ	-8.96	1.26	1.38
1	A	126	TYR	CE1-CZ	-8.21	1.27	1.38
1	B	126	TYR	CE1-CZ	-7.21	1.29	1.38
1	B	95	GLN	CD-NE2	-6.81	1.15	1.32

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	178	ARG	NE-CZ-NH2	21.32	130.96	120.30
1	B	219	ILE	CG1-CB-CG2	-9.85	89.72	111.40
1	D	178	ARG	NH1-CZ-NH2	-7.74	110.89	119.40
1	A	89	ARG	NE-CZ-NH2	7.70	124.15	120.30
1	A	178	ARG	NE-CZ-NH2	7.03	123.81	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2794	0	2683	61	0
1	B	2885	0	2824	65	1
1	C	2904	0	2861	63	1
1	D	2900	0	2871	68	0
2	A	31	0	0	4	0
2	B	25	0	0	1	0
2	C	32	0	0	3	0
2	D	15	0	0	0	0
All	All	11586	0	11239	221	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:ARG:NH1	1:B:65:PRO:HB3	1.69	1.06
1:A:89:ARG:HH11	1:B:65:PRO:HB3	1.24	1.00
1:A:58:CYS:HB3	1:A:89:ARG:HH21	1.38	0.89
1:C:19:TYR:HE1	1:D:132:TRP:CH2	1.92	0.88
1:A:89:ARG:HH11	1:B:65:PRO:CB	1.88	0.87

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:343:GLU:OE1	1:C:380:ARG:O[1_554]	2.02	0.18

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/413 (91%)	348 (92%)	26 (7%)	3 (1%)	19	43
1	B	391/413 (95%)	369 (94%)	21 (5%)	1 (0%)	41	66
1	C	386/413 (94%)	361 (94%)	23 (6%)	2 (0%)	29	54
1	D	387/413 (94%)	357 (92%)	28 (7%)	2 (0%)	29	54
All	All	1541/1652 (93%)	1435 (93%)	98 (6%)	8 (0%)	29	54

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	362	VAL
1	B	362	VAL
1	D	67	ILE
1	C	362	VAL
1	D	362	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/315 (86%)	252 (93%)	18 (7%)	16	37
1	B	285/315 (90%)	262 (92%)	23 (8%)	11	27
1	C	291/315 (92%)	272 (94%)	19 (6%)	17	38
1	D	293/315 (93%)	270 (92%)	23 (8%)	12	29
All	All	1139/1260 (90%)	1056 (93%)	83 (7%)	14	33

5 of 83 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	373	LEU
1	D	210	VAL
1	C	405	GLN
1	D	141	ILE
1	D	297	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN
1	B	181	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	391/413 (94%)	-0.15	7 (1%) 68 70	17, 42, 73, 103	0
1	B	397/413 (96%)	-0.18	7 (1%) 68 70	20, 41, 77, 90	0
1	C	396/413 (95%)	-0.13	7 (1%) 68 70	17, 43, 72, 94	0
1	D	395/413 (95%)	-0.05	12 (3%) 50 51	16, 44, 74, 97	0
All	All	1579/1652 (95%)	-0.13	33 (2%) 63 65	16, 43, 75, 103	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3	MET	4.5
1	C	342	GLY	3.9
1	A	249	THR	3.8
1	D	241	LEU	3.6
1	D	246	PRO	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.