



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 21, 2021 – 12:46 am BST

PDB ID : 6ZSM  
Title : Crystal structure of rsGCaMP double mutant Ile80His/Val116Ile in the ON state (non-illuminated)  
Authors : Janowski, R.; Fuenzalida-Werner, J.P.; Mishra, K.; Stiel, A.C.; Niessing, D.  
Deposited on : 2020-07-16  
Resolution : 1.95 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3729 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein, Green fluorescent protein, Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	399	3218	2022	545	636	15	0	9	2

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	36	MET	-	initiating methionine	UNP P42212
A	37	VAL	-	expression tag	UNP P42212
A	38	ASP	-	expression tag	UNP P42212
A	39	SER	-	expression tag	UNP P42212
A	40	SER	-	expression tag	UNP P42212
A	41	ARG	-	expression tag	UNP P42212
A	42	ARG	-	expression tag	UNP P42212
A	43	LYS	-	expression tag	UNP P42212
A	44	TRP	-	expression tag	UNP P42212
A	45	ASN	-	expression tag	UNP P42212
A	46	LYS	-	expression tag	UNP P42212
A	47	THR	-	expression tag	UNP P42212
A	48	GLY	-	expression tag	UNP P42212
A	49	HIS	-	expression tag	UNP P42212
A	50	ALA	-	expression tag	UNP P42212
A	51	VAL	-	expression tag	UNP P42212
A	52	ARG	-	expression tag	UNP P42212
A	53	ALA	-	expression tag	UNP P42212
A	54	ILE	-	expression tag	UNP P42212
A	55	GLY	-	expression tag	UNP P42212
A	56	ARG	-	expression tag	UNP P42212
A	57	LEU	-	expression tag	UNP P42212
A	58	SER	-	expression tag	UNP P42212
A	59	SER	-	expression tag	UNP P42212
A	60	LEU	-	expression tag	UNP P42212
A	61	GLU	-	expression tag	UNP P42212
A	66	LYS	MET	conflict	UNP P42212

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	ALA	VAL	conflict	UNP P42212
A	80	HIS	ILE	conflict	UNP P42212
A	88	GLY	SER	conflict	UNP P42212
A	93	TYR	ASP	conflict	UNP P42212
A	116	ILE	THR	conflict	UNP P42212
A	119	LYS	ALA	conflict	UNP P42212
A	144	LEU	HIS	conflict	UNP P42212
A	152	GLY	-	linker	UNP P42212
A	153	GLY	-	linker	UNP P42212
A	154	THR	-	linker	UNP P42212
A	155	GLY	-	linker	UNP P42212
A	156	GLY	-	linker	UNP P42212
A	157	SER	-	linker	UNP P42212
A	158	MET	-	linker	UNP P42212
A	159	VAL	-	linker	UNP P42212
A	?	-	PHE	deletion	UNP P42212
A	?	-	SER	deletion	UNP P42212
A	222	LEU	TYR	conflict	UNP P42212
A	224	PIA	GLY	conflict	UNP P42212
A	227	LEU	GLN	conflict	UNP P42212
A	251	ILE	VAL	conflict	UNP P42212
A	373	THR	ILE	conflict	UNP K4DIE3
A	381	TYR	ASP	conflict	UNP K4DIE3
A	452	LEU	-	expression tag	UNP K4DIE3
A	453	GLU	-	expression tag	UNP K4DIE3
A	454	HIS	-	expression tag	UNP K4DIE3
A	455	HIS	-	expression tag	UNP K4DIE3

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	5	Total Ca 5 5	0	0

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0
3	A	1	Total C O 3 1 2	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	444	Total O 444 444	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein, Green fluorescent protein, Calmodulin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.54Å 121.54Å 96.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.46 – 1.95 47.41 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.46-1.95) 100.0 (47.41-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.157 , 0.198 0.169 , 0.206	Depositor DCC
$R_{free}$ test set	2665 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.4	Xtrriage
Anisotropy	0.063	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3729	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EDO, PIA, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.02	14/3278 (0.4%)	1.05	9/4413 (0.2%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	269[A]	GLU	CD-OE2	8.73	1.35	1.25
1	A	269[B]	GLU	CD-OE2	8.73	1.35	1.25
1	A	287	ASP	CG-OD2	7.06	1.41	1.25
1	A	273	GLU	CD-OE2	6.53	1.32	1.25
1	A	192	GLU	CD-OE2	-6.29	1.18	1.25
1	A	390	GLU	CD-OE1	5.98	1.32	1.25
1	A	300	GLU	CD-OE2	-5.87	1.19	1.25
1	A	282	GLU	CD-OE1	5.42	1.31	1.25
1	A	135	GLU	CD-OE2	-5.41	1.19	1.25
1	A	386	GLU	CD-OE1	5.37	1.31	1.25
1	A	320	SER	CB-OG	5.29	1.49	1.42
1	A	334	GLU	CD-OE2	5.27	1.31	1.25
1	A	442	GLU	CD-OE1	5.26	1.31	1.25
1	A	163	GLU	CD-OE2	5.13	1.31	1.25

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	367	ASP	CB-CA-C	5.74	121.88	110.40
1	A	267	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	A	367	ASP	CB-CG-OD2	-5.25	113.58	118.30
1	A	421	ASP	N-CA-CB	-5.25	101.16	110.60
1	A	377	ARG	CG-CD-NE	-5.19	100.90	111.80
1	A	261	ASP	CB-CA-C	5.16	120.72	110.40
1	A	287	ASP	CA-CB-CG	5.15	124.72	113.40

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	306	GLN	N-CA-C	-5.07	97.30	111.00
1	A	340	ARG	NE-CZ-NH2	5.03	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3218	0	3138	36	0
2	A	5	0	0	0	0
3	A	42	0	14	0	0
4	A	20	0	30	3	0
5	A	444	0	0	9	0
All	All	3729	0	3182	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275[A]:ASP:OD1	5:A:603:HOH:O	1.73	1.06
1:A:128[A]:ARG:HH21	1:A:128[A]:ARG:HG3	1.28	0.96
1:A:128[A]:ARG:NH1	5:A:604:HOH:O	2.14	0.80
1:A:43:LYS:HD3	1:A:419:LEU:HD11	1.66	0.77
1:A:380:LYS:NZ	1:A:381:TYR:OH	2.20	0.75
1:A:126:GLU:OE2	1:A:128[A]:ARG:HG3	1.89	0.73
1:A:70:GLN:HG3	5:A:764:HOH:O	1.92	0.69
1:A:91:LEU:HD11	4:A:524:EDO:H11	1.76	0.67
1:A:418:LYS:O	1:A:419:LEU:HD23	1.94	0.66
1:A:367:ASP:CB	5:A:605:HOH:O	2.45	0.63
1:A:126:GLU:OE2	1:A:128[A]:ARG:NH2	2.33	0.62
1:A:128[A]:ARG:HG3	1:A:128[A]:ARG:NH2	2.04	0.60
1:A:224:PIA:N2	1:A:224:PIA:HD1	2.16	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71:LYS:O	1:A:99[A]:THR:HG23	2.05	0.56
1:A:448[A]:MET:HE2	4:A:521:EDO:H12	1.86	0.56
1:A:297:HIS:HD2	5:A:991:HOH:O	1.90	0.54
1:A:99[A]:THR:HG22	1:A:100:PRO:HD2	1.89	0.54
1:A:183:HIS:HE1	5:A:967:HOH:O	1.92	0.53
1:A:409:ARG:O	1:A:413:THR:HG23	2.10	0.51
1:A:305:ASP:OD1	1:A:380:LYS:HE3	2.13	0.48
1:A:105:PRO:HA	5:A:751:HOH:O	2.15	0.46
1:A:176:LEU:C	1:A:176:LEU:HD23	2.37	0.45
1:A:448[A]:MET:CE	4:A:521:EDO:H12	2.46	0.45
1:A:98:ASN:HA	1:A:251:ILE:O	2.17	0.45
1:A:367:ASP:HB3	5:A:605:HOH:O	2.12	0.45
1:A:64:TYR:O	1:A:76:ALA:HA	2.17	0.45
1:A:66:LYS:HE2	1:A:66:LYS:HB2	1.54	0.44
1:A:377:ARG:NH2	1:A:377:ARG:HG2	2.31	0.44
1:A:99[A]:THR:CG2	1:A:100:PRO:HD2	2.47	0.44
1:A:79:LYS:HB3	1:A:79:LYS:HE2	1.88	0.42
1:A:46:LYS:NZ	1:A:417:GLU:OE1	2.53	0.41
1:A:347:THR:OG1	1:A:350:GLU:HG3	2.20	0.41
1:A:63:VAL:O	1:A:113:TYR:HA	2.20	0.41
1:A:393:ARG:NH2	5:A:617:HOH:O	2.43	0.41
1:A:136:PHE:HA	1:A:198:GLY:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	401/418 (96%)	391 (98%)	10 (2%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	351/358 (98%)	339 (97%)	12 (3%)	37 25

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ASP
1	A	40	SER
1	A	42	ARG
1	A	66	LYS
1	A	127	LYS
1	A	128[A]	ARG
1	A	128[B]	ARG
1	A	210	LYS
1	A	304	ARG
1	A	374	MET
1	A	393	ARG
1	A	417	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	183	HIS
1	A	297	HIS
1	A	344	GLN
1	A	410	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PIA	A	224	1	21,21,22	4.60	5 (23%)	27,29,31	4.77	17 (62%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PIA	A	224	1	-	0/8/27/28	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	224	PIA	CA2-C2	-15.22	1.33	1.48
1	A	224	PIA	CB2-CA2	12.87	1.45	1.35
1	A	224	PIA	CA2-N2	3.58	1.46	1.38
1	A	224	PIA	O2-C2	3.58	1.30	1.23
1	A	224	PIA	CD1-CG2	-2.37	1.34	1.39

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	PIA	O2-C2-CA2	11.43	137.38	130.96
1	A	224	PIA	C2-N3-C1	9.42	112.73	107.97
1	A	224	PIA	CA1-C1-N2	7.34	133.44	124.05
1	A	224	PIA	CE1-CD1-CG2	-7.13	111.96	121.25
1	A	224	PIA	CA1-C1-N3	-6.56	116.88	124.75
1	A	224	PIA	CB2-CA2-C2	-6.54	114.47	122.28
1	A	224	PIA	CD2-CG2-CD1	5.83	126.26	117.64
1	A	224	PIA	CG2-CB2-CA2	-5.58	123.11	129.94
1	A	224	PIA	CA3-N3-C1	5.40	133.64	127.16

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	224	PIA	CA3-N3-C2	-4.47	113.55	123.80
1	A	224	PIA	CB2-CA2-N2	3.67	133.92	128.83
1	A	224	PIA	C2-CA2-N2	3.67	111.50	108.93
1	A	224	PIA	CD2-CE2-CZ	-3.61	115.92	119.88
1	A	224	PIA	CA2-N2-C1	-3.57	103.14	105.77
1	A	224	PIA	CD1-CE1-CZ	2.98	123.15	119.88
1	A	224	PIA	N3-C1-N2	-2.83	109.49	111.45
1	A	224	PIA	O2-C2-N3	-2.44	119.50	124.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	224	PIA	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 5 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FMT	A	518	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	519	-	0,2,2	-	-	0,1,1	-	-
4	EDO	A	520	-	3,3,3	0.34	0	2,2,2	0.55	0
3	FMT	A	511	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	515	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	517	-	0,2,2	-	-	0,1,1	-	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FMT	A	506	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	509	-	0,2,2	-	-	0,1,1	-	-
4	EDO	A	522	-	3,3,3	0.44	0	2,2,2	0.39	0
3	FMT	A	513	-	0,2,2	-	-	0,1,1	-	-
4	EDO	A	521	-	3,3,3	0.26	0	2,2,2	0.54	0
4	EDO	A	523	-	3,3,3	0.16	0	2,2,2	0.20	0
3	FMT	A	510	-	0,2,2	-	-	0,1,1	-	-
4	EDO	A	524	-	3,3,3	0.49	0	2,2,2	0.61	0
3	FMT	A	508	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	516	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	512	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	514	-	0,2,2	-	-	0,1,1	-	-
3	FMT	A	507	-	0,2,2	-	-	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	524	-	-	1/1/1/1	-
4	EDO	A	522	-	-	1/1/1/1	-
4	EDO	A	521	-	-	1/1/1/1	-
4	EDO	A	520	-	-	0/1/1/1	-
4	EDO	A	523	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	521	EDO	O1-C1-C2-O2
4	A	522	EDO	O1-C1-C2-O2
4	A	524	EDO	O1-C1-C2-O2
4	A	523	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	521	EDO	2	0
4	A	524	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	398/418 (95%)	0.21	24 (6%) <span style="border: 1px solid red; padding: 2px;">21</span> <span style="border: 1px solid red; padding: 2px;">30</span>	22, 33, 75, 161	8 (2%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	433	ILE	5.4
1	A	424	VAL	3.8
1	A	305	ASP	3.7
1	A	421	ASP	3.6
1	A	419	LEU	2.9
1	A	306	GLN	2.8
1	A	420	THR	2.8
1	A	435	GLY	2.8
1	A	422	GLU	2.7
1	A	431	ALA	2.7
1	A	159	VAL	2.6
1	A	425	ASP	2.6
1	A	437	GLY	2.4
1	A	418	LYS	2.4
1	A	432	ASP	2.4
1	A	409	ARG	2.3
1	A	450	ALA	2.3
1	A	303	THR	2.3
1	A	38	ASP	2.3
1	A	446	GLN	2.2
1	A	40	SER	2.2
1	A	429	ARG	2.1
1	A	406	ALA	2.1
1	A	402	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PIA	A	224	20/21	0.98	0.09	22,27,31,32	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	FMT	A	511	3/3	0.75	0.32	74,74,81,83	0
3	FMT	A	515	3/3	0.79	0.32	75,75,81,84	0
3	FMT	A	513	3/3	0.84	0.27	52,52,66,71	0
3	FMT	A	507	3/3	0.84	0.15	43,43,58,59	0
4	EDO	A	523	4/4	0.86	0.27	62,64,65,68	0
3	FMT	A	512	3/3	0.87	0.18	59,59,59,71	0
3	FMT	A	509	3/3	0.88	0.22	52,52,65,67	0
4	EDO	A	524	4/4	0.88	0.15	44,47,48,53	0
3	FMT	A	516	3/3	0.89	0.23	57,57,69,71	0
4	EDO	A	522	4/4	0.89	0.24	44,45,45,49	0
3	FMT	A	514	3/3	0.89	0.17	66,66,68,71	0
3	FMT	A	510	3/3	0.89	0.32	52,52,61,65	0
3	FMT	A	517	3/3	0.90	0.21	71,71,74,74	0
3	FMT	A	518	3/3	0.91	0.12	48,48,58,63	0
3	FMT	A	506	3/3	0.91	0.12	51,51,58,62	0
4	EDO	A	521	4/4	0.92	0.21	57,59,62,64	0
2	CA	A	505	1/1	0.93	0.11	84,84,84,84	0
3	FMT	A	519	3/3	0.93	0.21	50,50,62,62	0
3	FMT	A	508	3/3	0.95	0.18	47,47,50,54	0
4	EDO	A	520	4/4	0.95	0.13	36,49,51,54	0
2	CA	A	501	1/1	0.97	0.07	44,44,44,44	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	504	1/1	0.98	0.05	42,42,42,42	0
2	CA	A	503	1/1	1.00	0.11	29,29,29,29	0
2	CA	A	502	1/1	1.00	0.12	27,27,27,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.