



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2023 – 05:57 PM JST

PDB ID : 5ZX8
Title : Crystal structure of peptidyl-tRNA hydrolase from *Thermus thermophilus*
Authors : Matsumoto, A.; Uehara, U.; Shimizu, Y.; Ueda, T.; Uchiumi, T.; Ito, K.
Deposited on : 2018-05-18
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1050 (1.06-0.94)
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)
RSRZ outliers	127900	1023 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peptidyl-tRNA hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	186	1475	946	271	251	7	0	5	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q5SHZ2
A	-2	SER	-	expression tag	UNP Q5SHZ2
A	-1	HIS	-	expression tag	UNP Q5SHZ2
A	0	MET	-	expression tag	UNP Q5SHZ2

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: C₆H₅O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	13	6	7	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	A	1	Total	C	O	0	0
			8	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	120	Total	O	0	0
			120	120		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peptidyl-tRNA hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.45Å 53.92Å 58.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.45 – 1.00 23.44 – 1.00	Depositor EDS
% Data completeness (in resolution range)	92.2 (23.45-1.00) 92.3 (23.44-1.00)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.77 (at 1.00Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.129 , 0.143 0.138 , 0.149	Depositor DCC
R_{free} test set	3777 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	8.7	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 67.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1637	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.71	1/1525 (0.1%)	1.00	8/2060 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	166	GLU	CG-CD	5.38	1.60	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121	ARG	NE-CZ-NH1	12.49	126.55	120.30
1	A	121	ARG	NE-CZ-NH2	-7.91	116.35	120.30
1	A	34	ARG	NE-CZ-NH1	7.86	124.23	120.30
1	A	76	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	A	36	ARG	NE-CZ-NH1	6.55	123.58	120.30
1	A	34	ARG	NE-CZ-NH2	-6.26	117.17	120.30
1	A	142	ARG	NE-CZ-NH2	-6.13	117.23	120.30
1	A	23	PHE	CB-CG-CD1	5.00	124.30	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1475	0	1518	6	1
2	A	26	0	10	0	0
3	A	16	0	28	1	0
4	A	120	0	0	2	0
All	All	1637	0	1556	7	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:ARG:HG2	4:A:414:HOH:O	1.92	0.67
1:A:0:MET:SD	1:A:48:LEU:HG	2.37	0.64
3:A:204:MPD:H12	3:A:204:MPD:H52	1.86	0.56
1:A:43:GLU:HG2	1:A:48:LEU:HD23	1.94	0.49
1:A:96:GLY:HA2	1:A:115:ARG:O	2.14	0.47
1:A:148:VAL:CG2	1:A:182:LEU:HG	2.48	0.43
1:A:34:ARG:CG	4:A:318:HOH:O	2.68	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:GLU:OE1	1:A:174[A]:ARG:NH2[3_655]	1.86	0.34

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	189/187 (101%)	187 (99%)	2 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	152/147 (103%)	148 (97%)	4 (3%)	46 13

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ARG
1	A	19	HIS
1	A	34	ARG
1	A	98	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MPD	A	203	-	7,7,7	0.43	0	9,10,10	1.14	1 (11%)
3	MPD	A	204	-	7,7,7	0.76	0	9,10,10	0.82	0
2	FLC	A	202	-	12,12,12	3.11	4 (33%)	17,17,17	2.50	8 (47%)
2	FLC	A	201	-	12,12,12	3.08	6 (50%)	17,17,17	2.65	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	203	-	-	3/5/5/5	-
3	MPD	A	204	-	-	2/5/5/5	-
2	FLC	A	202	-	-	4/16/16/16	-
2	FLC	A	201	-	-	5/16/16/16	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	202	FLC	CB-CBC	-7.69	1.45	1.53
2	A	201	FLC	CA-CB	-6.15	1.46	1.53
2	A	201	FLC	CG-CB	-5.47	1.47	1.53
2	A	202	FLC	CG-CB	-5.43	1.47	1.53
2	A	201	FLC	OHB-CB	4.51	1.52	1.43
2	A	202	FLC	CA-CB	-3.43	1.49	1.53
2	A	201	FLC	OA2-CAC	-2.41	1.22	1.30
2	A	202	FLC	OHB-CB	2.28	1.47	1.43
2	A	201	FLC	OG1-CGC	2.14	1.29	1.22
2	A	201	FLC	OG2-CGC	-2.09	1.23	1.30

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	FLC	OB2-CBC-CB	6.69	124.67	113.05
2	A	201	FLC	OB1-CBC-CB	-5.30	114.74	122.25
2	A	202	FLC	CG-CB-CA	4.93	122.01	109.16
2	A	202	FLC	OHB-CB-CG	-4.17	99.63	109.40
2	A	202	FLC	OHB-CB-CA	-3.86	100.36	109.40
2	A	201	FLC	CG-CB-CA	3.65	118.69	109.16
2	A	201	FLC	OHB-CB-CG	-3.50	101.21	109.40
2	A	202	FLC	OHB-CB-CBC	-3.04	104.59	108.86
2	A	202	FLC	CB-CA-CAC	-2.70	107.29	113.81
2	A	202	FLC	OB2-CBC-CB	2.68	117.71	113.05
3	A	203	MPD	CM-C2-C1	-2.51	105.35	110.57
2	A	201	FLC	OHB-CB-CA	-2.49	103.57	109.40
2	A	202	FLC	CG-CB-CBC	2.46	115.39	110.11
2	A	201	FLC	CA-CB-CBC	2.43	115.32	110.11
2	A	202	FLC	OG2-CGC-CG	2.21	121.44	114.35

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	203	MPD	C1-C2-C3-C4
3	A	203	MPD	O2-C2-C3-C4
3	A	204	MPD	C2-C3-C4-O4
2	A	201	FLC	CA-CB-CG-CGC
3	A	203	MPD	CM-C2-C3-C4
2	A	201	FLC	CG-CB-CBC-OB2
2	A	202	FLC	CA-CB-CBC-OB1
2	A	202	FLC	CA-CB-CBC-OB2
2	A	202	FLC	CA-CB-CG-CGC
2	A	202	FLC	CAC-CA-CB-OHB
2	A	201	FLC	CG-CB-CBC-OB1
2	A	201	FLC	CB-CA-CAC-OA1
3	A	204	MPD	O2-C2-C3-C4
2	A	201	FLC	CB-CA-CAC-OA2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	204	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	186/187 (99%)	2.17	97 (52%) 0 1	9, 13, 25, 39	0

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	LEU	7.7
1	A	-3	GLY	6.0
1	A	-2	SER	4.5
1	A	21[A]	LEU	4.4
1	A	48	LEU	4.3
1	A	-1	HIS	4.2
1	A	80	VAL	3.9
1	A	122	LEU	3.9
1	A	108	ILE	3.8
1	A	79	VAL	3.7
1	A	78	LEU	3.7
1	A	4	VAL	3.6
1	A	5	VAL	3.5
1	A	1	MET	3.5
1	A	31	LEU	3.5
1	A	163	TRP	3.4
1	A	77	ILE	3.4
1	A	63	VAL	3.4
1	A	25	VAL	3.3
1	A	120	LEU	3.2
1	A	26	LEU	3.2
1	A	23	PHE	3.2
1	A	162	CYS	3.1
1	A	106	LEU	3.1
1	A	152	VAL	3.1
1	A	124	ILE	3.0
1	A	33	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	105	VAL	3.0
1	A	153	LEU	3.0
1	A	160	VAL	2.9
1	A	58	LEU	2.9
1	A	55	TYR	2.9
1	A	142	ARG	2.9
1	A	112	LEU	2.8
1	A	156	ALA	2.8
1	A	3	LEU	2.8
1	A	40	LEU	2.8
1	A	42	ALA	2.8
1	A	29	LEU	2.8
1	A	171	CYS	2.8
1	A	169	PRO	2.8
1	A	91	ILE	2.8
1	A	56	TYR	2.8
1	A	117	PHE	2.8
1	A	14	TYR	2.7
1	A	49	PHE	2.7
1	A	141	PHE	2.7
1	A	149	VAL	2.7
1	A	8	GLY	2.7
1	A	159	ALA	2.7
1	A	50	LEU	2.6
1	A	34	ARG	2.6
1	A	41	VAL	2.6
1	A	6	GLY	2.6
1	A	148	VAL	2.6
1	A	53	LEU	2.6
1	A	64	ALA	2.6
1	A	155	ALA	2.6
1	A	170	PRO	2.6
1	A	36	ARG	2.5
1	A	115	ARG	2.5
1	A	161	TRP	2.5
1	A	66	LEU	2.5
1	A	88	LEU	2.5
1	A	17	THR	2.4
1	A	85	ASP	2.4
1	A	72	ILE	2.4
1	A	167	GLY	2.4
1	A	178	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	93	PHE	2.4
1	A	164	VAL	2.3
1	A	46	GLY	2.3
1	A	59	THR	2.3
1	A	86	LEU	2.3
1	A	128	PRO	2.3
1	A	22	GLY	2.3
1	A	69	PHE	2.3
1	A	95	ALA	2.2
1	A	74	PRO	2.2
1	A	136	TYR	2.2
1	A	97	GLY	2.2
1	A	138	LEU	2.2
1	A	15	ALA	2.2
1	A	137	VAL	2.2
1	A	35	PRO	2.2
1	A	96	GLY	2.1
1	A	180	LEU	2.1
1	A	57	ASN	2.1
1	A	107[A]	SER	2.1
1	A	2	PHE	2.1
1	A	175	PHE	2.1
1	A	44	ALA	2.1
1	A	20	ASN	2.0
1	A	52	PRO	2.0
1	A	27	ASP	2.0
1	A	54	THR	2.0
1	A	176	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	A	203	8/8	0.77	0.17	20,25,28,40	0
3	MPD	A	204	8/8	0.80	0.17	19,24,28,33	0
2	FLC	A	202	13/13	0.86	0.21	19,21,25,26	0
2	FLC	A	201	13/13	0.93	0.15	15,16,19,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.