



Full wwPDB EM Validation Report ⓘ

Dec 18, 2022 – 07:01 am GMT

PDB ID : 7ZX7
EMDB ID : EMD-15006
Title : Structure of SNAPc containing Pol II pre-initiation complex bound to U1 snRNA promoter (CC)
Authors : Rengachari, S.; Schilbach, S.; Kaliyappan, T.; Gouge, J.; Zumer, K.; Schwarz, J.; Urlaub, H.; Dienemann, C.; Vannini, A.; Cramer, P.
Deposited on : 2022-05-20
Resolution : 3.40 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1970	72% 28%
2	B	1174	97% .
3	C	275	93% 7%
4	D	142	90% 10%
5	E	210	99% .
6	F	127	62% 38%
7	G	172	99% ..
8	H	150	99% .
9	I	125	91% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
10	J	67	96%
11	K	117	98%
12	L	58	76% 24%
13	M	316	77% 23%
14	N	96	54% 15% 31%
15	O	339	53% 47%
16	Q	517	26% 73%
17	R	249	89% 11%
18	T	96	55% 14% 31%
19	U	376	23% 77%
20	V	109	89% 11%
21	a	368	10% 58% 42%
22	b	411	89% 11%
23	c	1469	21% 79%
24	d	98	27% 53% 47%

2 Entry composition [i](#)

There are 26 unique types of molecules in this entry. The entry contains 49277 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1423	11274	7092	2016	2094	72	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1136	9076	5739	1597	1676	64	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	257	2059	1294	351	408	6	0	0

- Molecule 4 is a protein called RNA polymerase II subunit D.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	128	1050	656	178	212	4	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerase II subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	209	1721	1089	300	324	8	0	0

- Molecule 6 is a protein called DNA-directed RNA polymerase II subunit F.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	79	636	406	108	117	5	0	0

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	G	171	1351	875	219	249	8	0	0

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	H	148	1186	750	194	237	5	0	0

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	I	114	928	571	166	180	11	0	0

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	J	64	507	328	86	87	6	0	0

- Molecule 11 is a protein called RNA polymerase II subunit J.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	K	115	920	593	152	173	2	0	0

- Molecule 12 is a protein called RNA polymerase II subunit K.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	L	44	373	231	72	64	6	0	0

- Molecule 13 is a protein called Transcription initiation factor IIB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	M	242	1879	1182	332	349	16	0	0

- Molecule 14 is a DNA chain called Non-template strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	66	Total	C	N	O	P	0	0
			1369	648	258	397	66		

- Molecule 15 is a protein called TATA-box-binding protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	179	Total	C	N	O	S	0	0
			1422	923	251	241	7		

- Molecule 16 is a protein called General transcription factor IIF subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	Q	138	Total	C	N	O	S	0	0
			1138	719	208	208	3		

- Molecule 17 is a protein called General transcription factor IIF subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	R	222	Total	C	N	O	S	0	0
			1788	1127	320	338	3		

- Molecule 18 is a DNA chain called Template strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	T	66	Total	C	N	O	P	0	0
			1337	637	239	395	66		

- Molecule 19 is a protein called Transcription initiation factor IIA subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	U	88	Total	C	N	O	S	0	0
			734	470	124	136	4		

- Molecule 20 is a protein called Transcription initiation factor IIA subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	V	97	Total	C	N	O	S	0	0
			793	502	140	149	2		

- Molecule 21 is a protein called snRNA-activating protein complex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	a	215	1807	1165	313	319	10	0	0

- Molecule 22 is a protein called snRNA-activating protein complex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
22	b	365	2977	1890	510	557	20	0	0

- Molecule 23 is a protein called snRNA-activating protein complex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	c	303	2516	1574	453	479	10	0	0

- Molecule 24 is a protein called snRNA-activating protein complex subunit 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
24	d	52	424	259	81	81	3	0	0

- Molecule 25 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
25	A	2	Total	Zn	0
			2	2	
25	B	1	Total	Zn	0
			1	1	
25	C	1	Total	Zn	0
			1	1	
25	I	2	Total	Zn	0
			2	2	
25	J	1	Total	Zn	0
			1	1	
25	L	1	Total	Zn	0
			1	1	
25	M	1	Total	Zn	0
			1	1	
25	b	2	Total	Zn	0
			2	2	


- Molecule 26 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Lig-

and of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
26	A	1	1	1	0


- Molecule 3: DNA-directed RNA polymerase II subunit RPB3

Chain C:  93% 7%



- Molecule 4: RNA polymerase II subunit D

Chain D:  90% 10%




- Molecule 5: DNA-directed RNA polymerase II subunit E

Chain E:  99%




- Molecule 6: DNA-directed RNA polymerase II subunit F

Chain F:  62% 38%



- Molecule 7: DNA-directed RNA polymerase II subunit RPB7

Chain G:  99%





- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3


Chain H:  99%



- Molecule 9: DNA-directed RNA polymerase II subunit RPB9

Chain I:  91% 9%





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	47293	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	54.45	Depositor
Minimum defocus (nm)	300	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.193	Depositor
Minimum map value	-0.055	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.0183	Depositor
Map size (\AA)	419.99997, 419.99997, 419.99997	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.05, 1.05, 1.05	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/11479	0.41	0/15496
2	B	0.28	0/9257	0.42	0/12493
3	C	0.27	0/2102	0.43	0/2857
4	D	0.24	0/1064	0.35	0/1428
5	E	0.25	0/1752	0.40	0/2366
6	F	0.27	0/646	0.40	0/871
7	G	0.25	0/1382	0.40	0/1874
8	H	0.28	0/1207	0.46	0/1628
9	I	0.25	0/949	0.44	0/1284
10	J	0.29	0/516	0.42	0/696
11	K	0.27	0/939	0.42	0/1271
12	L	0.31	0/378	0.40	0/500
13	M	0.25	0/1909	0.42	0/2580
14	N	0.94	14/1538 (0.9%)	0.88	0/2376
15	O	0.25	0/1448	0.43	0/1948
16	Q	0.24	0/1167	0.39	0/1576
17	R	0.29	0/1817	0.45	0/2445
18	T	1.03	13/1496 (0.9%)	0.88	0/2302
19	U	0.24	0/747	0.43	0/1005
20	V	0.25	0/803	0.45	0/1088
21	a	0.24	0/1851	0.39	0/2493
22	b	0.24	0/3053	0.43	0/4131
23	c	0.24	0/2562	0.40	0/3440
24	d	0.24	0/425	0.46	0/564
All	All	0.35	27/50487 (0.1%)	0.46	0/68712

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	N	-7	DG	C1'-N9	-7.92	1.36	1.47
14	N	-9	DG	C1'-N9	-7.87	1.36	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	N	24	DA	C1'-N9	-7.56	1.36	1.47
18	T	-28	DA	C1'-N9	-7.54	1.36	1.47
18	T	-9	DA	C1'-N9	-7.51	1.36	1.47
18	T	5	DG	C1'-N9	-7.17	1.37	1.47
14	N	20	DG	C1'-N9	-7.01	1.37	1.47
18	T	-23	DG	C1'-N9	-6.99	1.37	1.47
18	T	-21	DG	C1'-N9	-6.92	1.37	1.47
14	N	34	DA	C1'-N9	-6.76	1.37	1.47
18	T	3	DA	C1'-N9	-6.74	1.37	1.47
14	N	25	DA	C1'-N9	-6.74	1.37	1.47
14	N	32	DA	C1'-N9	-6.72	1.37	1.47
14	N	-14	DA	C1'-N9	-6.66	1.38	1.47
14	N	-8	DA	C1'-N9	-6.47	1.38	1.47
14	N	-15	DA	C1'-N9	-6.44	1.38	1.47
18	T	-29	DA	C1'-N9	-6.36	1.38	1.47
14	N	38	DA	C1'-N9	-6.31	1.38	1.47
18	T	-39	DG	C1'-N9	-6.13	1.38	1.47
14	N	27	DG	C1'-N9	-6.11	1.38	1.47
14	N	0	DG	C1'-N9	-6.09	1.38	1.47
18	T	-10	DG	C1'-N9	-6.07	1.38	1.47
18	T	-15	DG	C1'-N9	-6.03	1.38	1.47
18	T	10	DA	C1'-N9	-5.87	1.39	1.47
18	T	-20	DC	C1'-N1	5.85	1.56	1.49
18	T	21	DA	C1'-N9	-5.79	1.39	1.47
14	N	22	DA	C1'-N9	-5.69	1.39	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1413/1970 (72%)	1377 (98%)	36 (2%)	0	100	100
2	B	1130/1174 (96%)	1098 (97%)	32 (3%)	0	100	100
3	C	253/275 (92%)	247 (98%)	6 (2%)	0	100	100
4	D	126/142 (89%)	124 (98%)	2 (2%)	0	100	100
5	E	207/210 (99%)	203 (98%)	4 (2%)	0	100	100
6	F	77/127 (61%)	74 (96%)	3 (4%)	0	100	100
7	G	169/172 (98%)	165 (98%)	4 (2%)	0	100	100
8	H	146/150 (97%)	143 (98%)	3 (2%)	0	100	100
9	I	112/125 (90%)	106 (95%)	6 (5%)	0	100	100
10	J	62/67 (92%)	62 (100%)	0	0	100	100
11	K	113/117 (97%)	112 (99%)	1 (1%)	0	100	100
12	L	42/58 (72%)	41 (98%)	1 (2%)	0	100	100
13	M	238/316 (75%)	231 (97%)	7 (3%)	0	100	100
15	O	177/339 (52%)	174 (98%)	3 (2%)	0	100	100
16	Q	134/517 (26%)	129 (96%)	5 (4%)	0	100	100
17	R	218/249 (88%)	211 (97%)	7 (3%)	0	100	100
19	U	84/376 (22%)	79 (94%)	5 (6%)	0	100	100
20	V	95/109 (87%)	87 (92%)	8 (8%)	0	100	100
21	a	211/368 (57%)	207 (98%)	4 (2%)	0	100	100
22	b	359/411 (87%)	336 (94%)	23 (6%)	0	100	100
23	c	299/1469 (20%)	288 (96%)	11 (4%)	0	100	100
24	d	50/98 (51%)	50 (100%)	0	0	100	100
All	All	5715/8839 (65%)	5544 (97%)	171 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1254/1749 (72%)	1253 (100%)	1 (0%)	93	98
2	B	994/1027 (97%)	991 (100%)	3 (0%)	92	97
3	C	234/252 (93%)	234 (100%)	0	100	100
4	D	118/126 (94%)	118 (100%)	0	100	100
5	E	191/192 (100%)	189 (99%)	2 (1%)	76	88
6	F	69/111 (62%)	69 (100%)	0	100	100
7	G	152/153 (99%)	151 (99%)	1 (1%)	84	92
8	H	129/131 (98%)	129 (100%)	0	100	100
9	I	103/112 (92%)	103 (100%)	0	100	100
10	J	53/56 (95%)	53 (100%)	0	100	100
11	K	104/106 (98%)	104 (100%)	0	100	100
12	L	41/55 (74%)	41 (100%)	0	100	100
13	M	206/268 (77%)	206 (100%)	0	100	100
15	O	154/293 (53%)	154 (100%)	0	100	100
16	Q	121/448 (27%)	120 (99%)	1 (1%)	81	91
17	R	196/218 (90%)	195 (100%)	1 (0%)	88	94
19	U	82/324 (25%)	82 (100%)	0	100	100
20	V	89/98 (91%)	89 (100%)	0	100	100
21	a	198/334 (59%)	197 (100%)	1 (0%)	88	94
22	b	323/356 (91%)	323 (100%)	0	100	100
23	c	270/1213 (22%)	270 (100%)	0	100	100
24	d	48/93 (52%)	48 (100%)	0	100	100
All	All	5129/7715 (66%)	5119 (100%)	10 (0%)	93	98

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	ARG
2	B	199	LYS
2	B	897	ARG
2	B	1131	ARG
5	E	52	ARG
5	E	162	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
7	G	81	LYS
16	Q	151	ARG
17	R	230	LYS
21	a	200	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-15006. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 218




Y Index: 208




Z Index: 243


6.3.2 Raw map



X Index: 218



Y Index: 208




Z Index: 243

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0183. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.5.1 emd_15006_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1079 nm^3 ; this corresponds to an approximate mass of 975 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.36	3.93	3.43
Unmasked-calculated*	4.30	8.32	4.43

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.30 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-15006 and PDB model 7ZX7. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0183 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0183).



















































9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0183) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9500	 0.2670
A	 0.9771	 0.3270
B	 0.9610	 0.3300
C	 0.9807	 0.3970
D	 0.8763	 0.1340
E	 0.9810	 0.2990
F	 0.9757	 0.3720
G	 0.9227	 0.1600
H	 0.9879	 0.3930
I	 0.9780	 0.2570
J	 0.9556	 0.3670
K	 0.9803	 0.4280
L	 0.9664	 0.2890
M	 0.9652	 0.3220
N	 0.9737	 0.2010
O	 0.9769	 0.2550
Q	 0.9315	 0.1400
R	 0.9441	 0.1430
T	 0.9723	 0.1930
U	 0.9583	 0.1760
V	 0.9703	 0.1960
a	 0.7914	 0.1270
b	 0.9577	 0.1220
c	 0.8779	 0.1260
d	 0.4587	 0.1120

