Summary of integrative structure determination of F1N6 fully-glycosylated model of mouse N-cadherin EC1-EC5 (PDB ID: 9A46, PDB-Dev ID: PDBDEV_00000227)

1. Model Composition	
1. Model Composition	- N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4))-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido
	galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]alpha-D-mannopyranose-(1-

6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain D (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain H (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)lalpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain M (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain I (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain G (18 residues)

Entry composition

- N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain N (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain P (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dqlucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain Q (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain R (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain J (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain K (18 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain O (18 residues) - Cadherin-2: Chain B (541 residues) - Cadherin-2: Chain A (541 residues) - N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]alpha-D-mannopyranose-(1-3)-[N-acetyl-alphaneuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-[N-acetyl-alpha-neuraminic acid-(2-3)-beta-Dgalactopyranose-(1-4)-2-acetamido-2-deoxy-beta-Dglucopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2deoxy-beta-D-glucopyranose-(1-4)-[alpha-Lfucopyranose-(1-6)]2-acetamido-2-deoxy-beta-Dglucopyranose: Chain L (18 residues) - SAS data, SASBDB: SASDT35 - Other, PDB: 3Q2W

Datasets used for modeling

2. Representation

Resolution	Atomic
Number of <u>rigid bodies</u> , <u>flexible units</u>	0, 18
Flexible units	- A: 1-541 - B: 1-541 - C: 1-18 - D: 1-18 - E: 1-18 - F: 1-18 - G: 1-18 - H: 1-18 - I: 1-18 - J: 1-18 - K: 1-18 - L: 1-18 - N: 1-18 - N: 1-18 - N: 1-18 - N: 1-18 - R: 1-18
Structural coverage (rigid bodies)	100%
3. Restraints	
Physical principles	Information about physical principles was not provided
Experimental data	- 1 unique SASRestraint: Assembly name: F1N6 fully-glycosylated model of mouse N-cadherin EC1-EC5 Fitting method: GASBOR Multi-state: True
4. Validation	
Number of ensembles	1
Number of models in ensembles	20
Number of deposited models	20
Model precision (uncertainty of models)	None, Å
Data quality	

Model quality: assessment of atomic segments

- Model-1: Clashscore = 24.7, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-2: Clashscore = 14.6, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-3: Clashscore = 11.75, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-4: Clashscore = 22.66, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-5: Clashscore = 14.31, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-6: Clashscore = 21.53, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-7: Clashscore = 18.52, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-8: Clashscore = 21.89, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-9: Clashscore = 10.03, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-10: Clashscore = 14.47, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-11: Clashscore = 13.68, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-12: Clashscore = 14.54, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-13: Clashscore = 15.86, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-14: Clashscore = 14.94, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-15: Clashscore = 10.44, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-16: Clashscore = 17.88, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-17: Clashscore = 10.99, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-18: Clashscore = 15.43, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-19: Clashscore = 11.92, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70
- Model-20: Clashscore = 14.66, Number of Ramachandran outliers = 6, Number of sidechain outliers = 70

Model quality: assessment of excluded volume	Not applicable
Fit to data used for modeling	Fit of model to information used to compute it has not been determined
Fit to data used for validation	Fit of model to information not used to compute it has not been determined
5. Methodology and Software	
1. <u>Method</u>	None
<u>Name</u>	Use GlycoSHIELD, the tool we have developed, to graft MD-simulated glycan ensemble onto the x-ray protein structure (PDB ID: 3Q2W).
Number of computed models	20
<u>Software</u>	- GlycoSHIELD (version Not available) - GASBOR (version Not available) - FoXSDock (version Not available)